

Table of Contents

Appendix G: Mortality.....	G-1
Figure G1a. Age-Adjusted All-Cause Mortality Rates for All Kansans, 1999-2013	G-2
Figure G1b. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013	G-5
Figure G1c. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013	G-8
Figure G1d. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013	G-11
Figure G1e. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-14
Figure G1f. Age-Adjusted All-Cause Mortality Rates for Hispanics in Kansas, 1999-2013	G-17
Figure G2a. Age-Adjusted Coronary Heart Disease Mortality Rates for All Kansans, 1999-2013	G-20
Figure G2b. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013	G-23
Figure G2c. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013.....	G-26
Figure G2d. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013.....	G-29
Figure G2e. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-32
Figure G2f. Age-Adjusted Coronary Heart Disease Mortality Rates for Hispanics in Kansas, 1999-2013	G-35
Figure G3a. Age-Adjusted Cancer Mortality Rates for All Kansans, 1999-2013.....	G-38
Figure G3b. Age-Adjusted Cancer Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013	G-41
Figure G3c. Age-Adjusted Cancer Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013	G-44
Figure G3d. Age-Adjusted Cancer Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013	G-47

Figure G3e. Age-Adjusted Cancer Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-50
Figure G3f. Age-Adjusted Cancer Mortality Rates for Hispanics in Kansas, 1999-2013....	G-53
Figure G4a. Age-Adjusted Diabetes Mortality Rates for All Kansans, 1999-2013	G-56
Figure G4b. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013	G-59
Figure G4c. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013	G-62
Figure G4d. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013	G-65
Figure G4e. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-68
Figure G4f. Age-Adjusted Diabetes Mortality Rates for Hispanics in Kansas, 1999-2013	G-71
Figure G5a. Age-Adjusted Unintentional Injury Mortality Rates for All Kansans, 1999-2013	G-74
Figure G5b. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013.....	G-77
Figure G5c. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013.....	G-80
Figure G5d. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013.....	G-83
Figure G5e. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-86
Figure G5f. Age-Adjusted Unintentional Injury Mortality Rates for Hispanics in Kansas, 1999-2013	G-89
Figure G6a. Age-Adjusted Homicide Mortality Rates for All Kansans, 1999-2013	G-92
Figure G6b. Age-Adjusted Homicide Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013	G-95
Figure G6c. Age-Adjusted Homicide Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013	G-98
Figure G6d. Age-Adjusted Homicide Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013	G-101

Figure G6e. Age-Adjusted Homicide Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-104
Figure G6f. Age-Adjusted Homicide Mortality Rates for Hispanics in Kansas, 1999-2013	G-107
Figure G7a. Age-Adjusted Suicide Mortality Rates for All Kansans, 1999-2013.....	G-110
Figure G7b. Age-Adjusted Suicide Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013	G-113
Figure G7c. Age-Adjusted Suicide Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013	G-116
Figure G7d. Age-Adjusted Suicide Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013	G-119
Figure G7e. Age-Adjusted Suicide Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-122
Figure G7f. Age-Adjusted Suicide Mortality Rates for Hispanics in Kansas, 1999-2013	G-125
Figure G8a. All-Cause Mortality Age-Adjusted YPLL* Rates for All Kansans, 1999-2013	G-128
Figure G8b. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013	G-131
Figure G8c. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013.....	G-134
Figure G8d. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013	G-137
Figure G8e. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-140
Figure G8f. All-Cause Mortality Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013	G-143
Figure G9a. Coronary Heart Disease Age-Adjusted YPLL* Rates for All Kansans, 1999-2013	G-146
Figure G9b. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013.....	G-149
Figure G9c. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013.....	G-152

Figure G9d. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013.....	G-155
Figure G9e. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-158
Figure G9f. Coronary Heart Disease Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013	G-161
Figure G10a. Cancer Age-Adjusted YPLL* Rates for All Kansans, 1999-2013	G-164
Figure G10b. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013	G-167
Figure G10c. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013	G-170
Figure G10d. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013	G-173
Figure G10e. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-176
Figure G10f. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-179
Figure G11a. Diabetes Age-Adjusted YPLL* Rates for All Kansans, 1999-2013.....	G-182
Figure G11b. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013	G-185
Figure G11c. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013	G-188
Figure G11d. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013	G-191
Figure G11e. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013.....	G-194
Figure G11f. Diabetes Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013	G-197
Figure G12a. Unintentional Injury Age-Adjusted YPLL* Rates for All Kansans, 1999-2013	G-200
Figure G12b. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013	G-203

Figure G12c. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013 G-206

Figure G12d. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013 G-209

Figure G12e. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013 G-212

Figure G12f. Unintentional Injury Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013 G-215

Appendix G: Mortality

Please note: “All Kansans” in the following figures refers to all Kansans in the subpopulation described in the title of the figures.

Figure G1a. Age-Adjusted All-Cause Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	123,040	841	122,194	798	123,968	759
Frontier	8,648	816	8,013	796	7,452	769
Rural	15,876	808	15,124	800	14,414	781
Dense Rural	24,346	862	23,588	830	23,376	808
Semi-Urban	19,992	841	20,091	812	20,511	786
Urban	54,166	849	55,326	782	58,204	732
Allen	935	917	861	901	933	963
Anderson	507	784	470	744	497	791
Atchison	928	870	927	884	887	844
Barber	345	834	353	868	293	806
Barton	1,612	825	1,536	788	1,498	790
Bourbon	1,009	910	953	909	897	852
Brown	678	798	729	941	645	895
Butler	2,644	853	2,757	822	3,037	834
Chase	215	967	163	729	162	710
Chautauqua	348	833	327	909	282	866
Cherokee	1,312	974	1,397	1,046	1,340	1,017
Cheyenne	198	708	215	757	202	677
Clark	146	676	158	746	161	746
Clay	533	710	575	752	529	714
Cloud	809	798	747	836	666	812
Coffey	526	839	538	917	512	867
Comanche	177	810	176	1,006	152	926
Cowley	2,224	957	2,071	896	2,135	918
Crawford	2,300	941	2,212	932	2,035	868
Decatur	261	756	264	765	256	840
Dickinson	1,244	863	1,113	758	1,175	792
Doniphan	440	831	389	746	388	711
Douglas	2,675	773	2,845	709	3,100	672
Edwards	234	876	203	797	197	874
Elk	298	972	238	865	215	856
Ellis	1,227	777	1,216	757	1,206	705
Ellsworth	429	771	399	745	417	796
Finney	1,041	831	936	700	995	692
Ford	1,361	917	1,148	769	1,187	760
Franklin	1,179	815	1,195	829	1,226	827
Geary	1,036	961	938	854	1,015	842
Gove	207	719	185	690	200	765

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1a. Age-Adjusted All-Cause Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	197	757	198	807	166	720
Grant	264	861	269	819	251	761
Gray	268	790	261	780	237	715
Greeley	88	871	76	730	75	689
Greenwood	646	975	547	901	469	870
Hamilton	150	817	151	884	151	1,001
Harper	523	866	513	946	461	896
Harvey	1,758	763	1,798	732	1,904	727
Haskell	156	863	159	819	132	667
Hodgeman	113	704	123	774	91	564
Jackson	667	866	656	832	635	747
Jefferson	789	785	863	799	862	744
Jewell	295	814	237	732	205	658
Johnson	13,913	702	15,523	654	17,235	623
Kearny	204	1,073	172	810	217	976
Kingman	568	818	535	770	483	738
Kiowa	193	759	156	713	139	689
Labette	1,501	955	1,434	959	1,311	900
Lane	150	870	134	906	113	760
Leavenworth	2,597	918	2,610	826	2,762	753
Lincoln	289	908	238	790	202	716
Linn	573	867	532	787	571	827
Logan	199	839	177	734	168	708
Lyon	1,450	799	1,471	805	1,418	778
McPherson	1,722	777	1,785	761	1,823	759
Marion	910	769	885	777	798	716
Marshall	739	753	658	729	644	710
Meade	250	759	233	705	251	739
Miami	1,257	875	1,217	757	1,285	716
Mitchell	483	760	481	802	473	844
Montgomery	2,376	925	2,368	958	2,180	893
Morris	396	797	359	724	385	729
Morton	143	745	187	866	190	815
Nemaha	742	710	722	758	645	689
Neosho	1,022	874	1,011	851	982	847
Ness	243	715	252	809	210	716
Norton	361	741	359	773	300	689
Osage	910	841	912	855	897	834

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1a. Age-Adjusted All-Cause Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	371	797	329	779	286	729
Ottawa	419	908	353	796	343	844
Pawnee	454	846	389	726	375	738
Phillips	466	881	372	741	375	821
Pottawatomie	847	832	841	777	794	688
Pratt	601	827	607	799	592	795
Rawlins	205	663	221	837	181	693
Reno	3,440	812	3,426	782	3,643	795
Republic	521	870	372	662	437	803
Rice	651	843	656	898	639	895
Riley	1,554	748	1,502	659	1,602	641
Rooks	375	808	345	814	319	759
Rush	288	820	256	784	231	717
Russell	515	773	540	854	469	747
Saline	2,448	828	2,678	849	2,584	767
Scott	276	773	263	778	300	841
Sedgwick	18,468	886	19,097	841	19,983	791
Seward	702	804	683	776	659	745
Shawnee	8,697	935	8,216	826	8,452	784
Sheridan	184	831	148	675	153	685
Sherman	361	835	373	851	375	868
Smith	374	747	330	736	307	734
Stafford	341	842	309	861	295	795
Stanton	99	813	96	697	85	597
Stevens	239	801	234	724	228	683
Sumner	1,403	874	1,385	888	1,354	860
Thomas	373	754	418	826	364	693
Trego	236	776	240	794	232	816
Wabaunsee	352	868	329	791	347	799
Wallace	107	895	83	629	90	681
Washington	496	793	424	742	374	681
Wichita	118	693	120	747	131	804
Wilson	740	962	682	911	651	939
Woodson	308	929	294	944	274	997
Wyandotte	7,816	1,078	7,035	998	6,672	908

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1b. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	113,578	835	110,829	787	111,337	752
Frontier	8,483	812	7,818	796	7,203	767
Rural	15,641	812	14,713	795	13,946	778
Dense Rural	23,190	866	22,150	830	21,752	814
Semi-Urban	19,021	843	18,798	805	18,985	776
Urban	47,232	836	47,319	763	49,444	717
Allen	909	919	818	884	899	966
Anderson	504	785	465	743	484	783
Atchison	866	858	865	869	830	839
Barber	339	832	347	860	288	807
Barton	1,558	821	1,473	788	1,417	794
Bourbon	974	909	907	891	873	863
Brown	640	792	658	902	566	841
Butler	2,593	857	2,637	810	2,866	814
Chase	212	965	160	730	155	684
Chautauqua	341	836	318	917	266	858
Cherokee	1,266	972	1,338	1,035	1,299	1,034
Cheyenne	196	702	214	764	201	692
Clark	144	668	155	747	156	746
Clay	530	711	569	748	523	715
Cloud	808	802	743	841	658	813
Coffey	517	827	531	916	506	873
Comanche	176	807	174	999	148	892
Cowley	2,095	948	1,936	889	1,930	891
Crawford	2,266	947	2,163	936	1,969	866
Decatur	260	758	263	763	255	845
Dickinson	1,212	858	1,073	744	1,141	786
Doniphan	419	815	373	738	369	701
Douglas	2,487	768	2,614	700	2,871	674
Edwards	225	887	199	836	179	882
Elk	289	930	234	870	214	874
Ellis	1,220	783	1,190	753	1,180	705
Ellsworth	422	782	389	753	407	811
Finney	842	858	727	719	749	743
Ford	1,242	970	1,006	821	980	801
Franklin	1,153	817	1,131	805	1,169	814
Geary	802	950	709	835	757	854
Gove	207	722	184	692	199	764

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1b. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	183	716	183	776	146	662
Grant	239	906	224	821	199	748
Gray	264	801	248	765	231	732
Greeley	85	837	74	743	70	671
Greenwood	642	978	541	905	454	863
Hamilton	140	791	143	877	143	1,035
Harper	521	871	505	945	445	876
Harvey	1,682	767	1,710	737	1,795	727
Haskell	145	896	139	815	113	640
Hodgeman	112	708	122	784	90	574
Jackson	608	825	592	791	565	709
Jefferson	780	791	838	789	846	746
Jewell	295	816	236	734	200	640
Johnson	13,379	706	14,602	654	16,122	624
Kearny	179	1,048	152	816	197	1,036
Kingman	561	814	531	775	478	743
Kiowa	190	746	154	715	137	691
Labette	1,424	962	1,336	953	1,220	901
Lane	148	855	134	924	110	732
Leavenworth	2,348	922	2,340	824	2,445	744
Lincoln	289	915	235	788	201	728
Linn	567	868	521	781	563	832
Logan	198	849	170	717	166	712
Lyon	1,375	817	1,363	815	1,292	794
McPherson	1,707	779	1,753	759	1,781	754
Marion	900	770	868	767	774	702
Marshall	737	757	651	727	636	702
Meade	237	745	226	703	229	689
Miami	1,212	867	1,167	747	1,243	716
Mitchell	481	759	478	799	466	834
Montgomery	2,177	922	2,138	937	1,983	889
Morris	390	797	352	724	379	734
Morton	140	765	168	810	173	803
Nemaha	738	710	709	753	641	691
Neosho	991	872	978	851	937	838
Ness	243	718	250	815	208	718
Norton	357	748	352	777	295	695
Osage	900	842	896	854	874	833

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1b. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	369	797	328	781	284	731
Ottawa	418	916	345	787	334	837
Pawnee	432	841	374	737	355	757
Phillips	466	887	370	748	372	828
Pottawatomie	833	834	825	779	766	682
Pratt	591	829	583	781	562	774
Rawlins	204	662	218	842	177	698
Reno	3,310	818	3,235	776	3,368	775
Republic	520	873	368	660	432	804
Rice	631	842	630	896	619	914
Riley	1,481	754	1,406	660	1,459	631
Rooks	373	810	340	812	317	764
Rush	285	811	254	786	228	719
Russell	511	774	532	853	461	748
Saline	2,345	836	2,502	842	2,376	757
Scott	270	764	253	769	293	877
Sedgwick	16,115	879	16,061	819	16,553	772
Seward	600	860	544	823	504	809
Shawnee	7,631	919	7,100	806	7,303	771
Sheridan	184	836	148	682	148	649
Sherman	348	825	354	842	357	873
Smith	374	749	329	737	306	740
Stafford	340	856	302	869	289	808
Stanton	91	844	87	724	74	619
Stevens	227	832	217	746	203	669
Sumner	1,369	882	1,319	880	1,300	865
Thomas	373	760	414	829	356	688
Trego	235	782	236	790	231	819
Wabaunsee	345	867	321	788	330	778
Wallace	103	868	82	641	89	678
Washington	496	798	420	739	371	680
Wichita	107	675	104	693	118	820
Wilson	733	967	663	898	625	919
Woodson	307	941	290	940	269	995
Wyandotte	5,272	1,082	4,602	1,027	4,150	935

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1c. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	6,255	1,155	6,238	1,078	6,135	936
Frontier	29	1,224	23	845	25	871
Rural	52	756	75	1,110	50	610
Dense Rural	354	1,083	373	1,145	338	907
Semi-Urban	597	1,092	635	1,200	602	989
Urban	5,223	1,178	5,128	1,063	5,118	941
Allen	17	1,247	19	1,600	12	893
Anderson	-	-	-	-	-	-
Atchison	49	1,175	42	1,001	35	814
Barber	-	-	-	-	-	-
Barton	12	693	14	779	10	581
Bourbon	29	1,050	35	1,699	14	580
Brown	-	-	19	1,281	11	762
Butler	18	961	27	1,029	26	869
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	12	1,429	15	2,287	7	1,183
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	47	1,109	34	810	65	1,347
Crawford	23	993	27	1,412	29	1,469
Decatur	-	-	-	-	-	-
Dickinson	9	1,388	6	1,012	6	690
Doniphan	15	1,864	6	747	6	595
Douglas	131	1,282	100	931	107	928
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	7	1,777	-	-
Ellsworth	-	-	-	-	-	-
Finney	9	772	8	502	11	509
Ford	12	1,103	13	909	13	818
Franklin	7	2,327	14	1,512	6	399
Geary	172	1,186	164	1,208	165	915
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1c. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	13	3,504	12	1,544	15	1,888
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	21	748	24	916	21	847
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	6	8,935	-	-	6	2,681
Jefferson	6	682	10	1,988	9	1,981
Jewell	-	-	-	-	-	-
Johnson	269	1,026	372	841	425	676
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	63	1,307	63	1,331	52	973
Lane	-	-	-	-	-	-
Leavenworth	217	1,057	208	999	218	986
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	18	728	25	1,250	23	960
McPherson	-	-	6	909	8	713
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	36	1,743	36	1,647	24	956
Mitchell	-	-	-	-	-	-
Montgomery	151	1,331	153	1,630	129	1,375
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	6	2,745	-	-
Neosho	11	1,492	13	1,993	13	1,692
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1c. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	12	987	12	1,074	7	460
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	6	1,833	-	-	6	1,080
Rawlins	-	-	-	-	-	-
Reno	65	1,047	61	1,023	68	1,047
Republic	-	-	-	-	-	-
Rice	-	-	6	2,158	-	-
Riley	35	701	45	853	67	791
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	69	1,063	84	1,292	67	877
Scott	-	-	-	-	-	-
Sedgwick	1,675	1,182	1,777	1,123	1,824	1,004
Seward	26	1,017	34	1,533	32	1,043
Shawnee	745	1,281	713	1,147	673	1,016
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	18	2,152	8	802
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	2,186	1,214	1,958	1,088	1,871	975

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1d. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	663	979	839	1,074	929	1,024
Frontier	28	1,270	33	1,118	41	1,223
Rural	67	1,375	92	1,435	111	1,332
Dense Rural	190	1,261	206	1,181	224	1,049
Semi-Urban	76	640	114	925	110	742
Urban	302	892	393	1,023	443	1,059
Allen	-	-	9	2,267	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	6	2,848	6	3,336
Barber	-	-	-	-	-	-
Barton	12	7,467	10	3,508	11	3,407
Bourbon	-	-	-	-	-	-
Brown	33	1,255	40	1,338	57	1,512
Butler	11	590	26	1,748	17	883
Chase	-	-	-	-	-	-
Chautauqua	-	-	6	765	14	1,745
Cherokee	28	1,168	18	708	16	426
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	37	1,632	28	1,068	57	2,476
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	9	6,549	-	-	-	-
Doniphan	-	-	8	2,693	8	3,304
Douglas	33	972	47	1,236	32	716
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	17	2,688	12	1,402
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1d. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	6	905	8	1,073
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	49	1,465	52	1,278	54	1,128
Jefferson	-	-	6	758	-	-
Jewell	-	-	-	-	-	-
Johnson	37	1,184	50	932	52	953
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	8	481	10	547	12	578
Lane	-	-	-	-	-	-
Leavenworth	7	539	10	778	13	680
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	6	5,598	9	3,059	11	1,803
McPherson	-	-	10	7,343	-	-
Marion	-	-	-	-	7	3,654
Marshall	-	-	-	-	-	-
Meade	8	7,654	-	-	6	2,965
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	19	440	21	471	20	388
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	7	1,430	8	1,814
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	6	1,372	8	992

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1d. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	9	10,294	6	11,396
Rawlins	-	-	-	-	-	-
Reno	12	1,004	19	1,602	21	1,455
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	7	1,452	9	2,507	6	1,088
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	10	1,398	9	981	16	1,562
Scott	-	-	-	-	-	-
Sedgwick	126	845	169	1,089	218	1,385
Seward	-	-	-	-	-	-
Shawnee	71	1,180	75	1,034	88	1,001
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	8	1,331	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	28	632	42	1,089	40	943

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1e. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	417	386	598	408	769	405
Frontier	-	-	-	-	-	-
Rural	8	638	10	423	21	587
Dense Rural	48	339	61	587	65	395
Semi-Urban	70	519	70	440	115	545
Urban	289	365	454	388	563	383
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	7	733
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	7	900	7	788	12	618
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	17	287	28	444
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	11	198	9	423	14	499
Ford	-	-	6	654	7	348
Franklin	-	-	-	-	-	-
Geary	39	823	29	571	44	681
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1e. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	8	669
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	72	257	149	360	171	324
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	10	301	13	363	19	390
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	7	515	11	530
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1e. Age-Adjusted All-Cause Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	6	546
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	10	495	9	173	16	334
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	11	398	24	577
Scott	-	-	-	-	-	-
Sedgwick	150	465	217	428	250	393
Seward	-	-	9	931	-	-
Shawnee	14	237	28	403	29	362
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	40	628	30	371	66	674

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1f. Age-Adjusted All-Cause Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	2,079	582	2,688	550	3,335	541
Frontier	103	824	105	652	134	608
Rural	105	525	170	543	197	555
Dense Rural	555	601	625	526	784	536
Semi-Urban	221	543	314	585	389	549
Urban	1,095	578	1,471	548	1,830	536
Allen	-	-	13	939	12	869
Anderson	-	-	-	-	6	1,370
Atchison	10	713	11	783	8	445
Barber	-	-	-	-	-	-
Barton	29	788	29	502	32	484
Bourbon	-	-	-	-	-	-
Brown	-	-	7	572	6	548
Butler	20	586	26	585	35	512
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	9	1,047	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	38	922	39	708	45	666
Crawford	-	-	10	416	15	441
Decatur	-	-	-	-	-	-
Dickinson	10	735	9	478	11	390
Doniphan	-	-	-	-	-	-
Douglas	20	516	36	396	35	344
Edwards	6	507	-	-	15	807
Elk	8	2,906	-	-	-	-
Ellis	-	-	8	886	12	742
Ellsworth	7	778	8	626	7	425
Finney	178	692	170	563	205	555
Ford	101	538	113	489	172	549
Franklin	14	806	13	668	15	694
Geary	17	485	26	514	25	300
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1f. Age-Adjusted All-Cause Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	25	657	40	721	48	958
Gray	-	-	11	628	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	9	1,360	6	2,411	7	465
Harper	-	-	-	-	8	1,130
Harvey	48	721	41	455	62	528
Haskell	9	517	17	1,035	17	687
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	152	475	241	473	290	436
Kearny	24	870	15	653	17	480
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	6	261	18	805	16	568
Lane	-	-	-	-	-	-
Leavenworth	15	350	26	497	42	546
Lincoln	-	-	-	-	-	-
Linn	-	-	7	1,966	-	-
Logan	-	-	-	-	-	-
Lyon	47	478	59	468	68	471
McPherson	-	-	8	449	17	579
Marion	-	-	7	924	6	442
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	12	1,545
Miami	8	917	6	314	7	310
Mitchell	-	-	-	-	-	-
Montgomery	25	695	34	808	31	670
Morris	-	-	-	-	-	-
Morton	-	-	14	1,419	13	980
Nemaha	-	-	-	-	-	-
Neosho	15	734	10	381	17	618
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	7	640	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G1f. Age-Adjusted All-Cause Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	8	1,066	-	-	9	513
Phillips	-	-	-	-	-	-
Pottawatomie	6	762	-	-	12	886
Pratt	-	-	8	665	12	627
Rawlins	-	-	-	-	-	-
Reno	48	484	79	653	98	631
Republic	-	-	-	-	-	-
Rice	12	676	15	704	15	620
Riley	20	541	18	418	27	372
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	19	318	61	777	74	691
Scott	6	1,139	8	1,416	-	-
Sedgwick	392	575	568	587	740	586
Seward	69	519	85	353	106	503
Shawnee	230	713	234	576	293	618
Sheridan	-	-	-	-	-	-
Sherman	12	2,572	16	1,475	10	706
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	7	330	11	716
Stevens	9	613	12	300	15	370
Sumner	22	601	26	716	28	613
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	11	1,267	15	1,132	9	583
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	286	577	366	568	430	525

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2a. Age-Adjusted Coronary Heart Disease Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	21,768	147	17,687	114	15,275	92
Frontier	1,941	170	1,426	130	1,110	104
Rural	2,934	143	2,408	122	1,900	98
Dense Rural	4,707	162	3,712	127	3,115	105
Semi-Urban	3,739	155	2,956	117	2,638	99
Urban	8,445	134	7,174	102	6,509	82
Allen	212	196	149	148	124	119
Anderson	108	157	93	133	55	84
Atchison	150	136	110	105	103	94
Barber	72	162	67	150	52	123
Barton	290	145	286	144	230	119
Bourbon	206	178	142	131	124	115
Brown	118	138	104	138	96	125
Butler	428	137	379	112	403	109
Chase	36	158	22	88	17	74
Chautauqua	83	187	51	127	42	114
Cherokee	384	278	272	196	216	159
Cheyenne	31	107	30	102	24	62
Clark	39	170	32	131	28	134
Clay	70	83	57	73	45	58
Cloud	169	153	123	125	80	96
Coffey	87	137	55	90	82	132
Comanche	26	106	38	187	30	177
Cowley	363	151	316	132	331	138
Crawford	622	245	413	169	340	141
Decatur	52	133	62	173	27	70
Dickinson	266	173	191	127	182	120
Doniphan	98	184	65	123	64	116
Douglas	390	115	314	79	313	69
Edwards	46	148	33	122	40	164
Elk	74	218	47	154	44	160
Ellis	206	125	171	104	148	84
Ellsworth	82	137	83	147	70	129
Finney	204	177	127	99	118	83
Ford	238	163	184	124	158	102
Franklin	215	143	175	120	133	87
Geary	164	160	122	116	148	131
Gove	44	134	29	98	31	106

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2a. Age-Adjusted Coronary Heart Disease Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	58	202	49	176	41	162
Grant	49	164	57	178	45	141
Gray	34	102	43	130	24	68
Greeley	18	172	9	81	10	74
Greenwood	163	228	101	153	78	134
Hamilton	19	106	29	165	32	208
Harper	103	166	87	156	45	81
Harvey	297	125	252	95	256	91
Haskell	46	261	42	224	28	143
Hodgeman	13	81	16	87	13	78
Jackson	123	155	94	115	76	87
Jefferson	130	128	125	113	126	105
Jewell	58	156	40	114	27	80
Johnson	2,008	102	1,812	76	1,764	63
Kearny	48	260	29	132	27	121
Kingman	117	155	122	166	75	107
Kiowa	35	131	19	83	11	48
Labette	308	189	225	143	145	93
Lane	24	132	22	154	22	158
Leavenworth	492	177	466	147	472	128
Lincoln	106	286	61	178	38	114
Linn	129	186	90	130	91	131
Logan	41	166	24	92	20	73
Lyon	241	134	178	99	142	76
McPherson	338	144	285	115	224	92
Marion	130	106	116	93	72	65
Marshall	157	148	108	114	111	109
Meade	52	147	39	115	36	93
Miami	296	206	164	102	223	122
Mitchell	91	135	71	113	62	105
Montgomery	444	164	406	154	295	111
Morris	70	142	45	78	55	97
Morton	44	227	46	202	22	87
Nemaha	115	115	96	103	74	72
Neosho	198	160	188	149	126	106
Ness	50	136	36	117	27	80
Norton	48	93	53	100	28	57
Osage	162	145	132	120	92	81

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2a. Age-Adjusted Coronary Heart Disease Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	135	243	76	152	72	151
Ottawa	102	212	38	86	37	90
Pawnee	78	140	57	103	43	78
Phillips	76	127	46	94	26	50
Pottawatomie	157	153	146	135	110	95
Pratt	84	109	100	121	68	89
Rawlins	47	140	47	161	26	78
Reno	535	123	424	93	395	86
Republic	82	146	50	81	53	97
Rice	106	137	95	121	92	129
Riley	260	127	207	91	142	60
Rooks	73	146	41	90	37	80
Rush	83	220	55	150	31	87
Russell	123	168	121	176	80	121
Saline	478	161	414	129	303	89
Scott	44	128	30	85	50	134
Sedgwick	2,918	142	2,521	111	2,333	92
Seward	143	171	84	100	87	103
Shawnee	1,572	167	1,233	123	922	84
Sheridan	35	154	29	129	25	113
Sherman	91	204	62	134	42	94
Smith	62	117	46	94	39	91
Stafford	85	196	53	139	35	84
Stanton	15	126	11	76	9	56
Stevens	51	173	42	125	30	84
Sumner	290	171	242	150	189	114
Thomas	72	140	56	113	43	75
Trego	42	130	45	132	29	93
Wabaunsee	76	185	50	116	39	92
Wallace	24	193	13	74	13	89
Washington	97	145	78	121	50	86
Wichita	17	98	17	86	13	75
Wilson	168	206	164	204	122	159
Woodson	52	167	36	112	29	97
Wyandotte	1,065	149	828	120	705	98

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2b. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	20,518	148	16,387	114	13,984	92
Frontier	1,923	171	1,386	129	1,081	104
Rural	2,886	143	2,356	121	1,851	98
Dense Rural	4,544	163	3,548	128	2,925	105
Semi-Urban	3,603	156	2,806	117	2,483	99
Urban	7,560	134	6,282	101	5,642	81
Allen	207	197	142	145	120	119
Anderson	108	158	92	132	54	83
Atchison	138	131	103	103	98	94
Barber	71	163	67	152	51	121
Barton	282	144	278	145	216	118
Bourbon	202	181	132	124	119	115
Brown	108	132	91	129	86	122
Butler	421	137	366	111	389	108
Chase	36	160	21	83	17	75
Chautauqua	83	191	51	131	40	113
Cherokee	373	278	263	194	210	160
Cheyenne	30	104	30	103	24	63
Clark	38	166	31	129	28	138
Clay	70	83	55	71	45	59
Cloud	169	154	121	122	79	96
Coffey	84	132	54	90	80	130
Comanche	26	107	38	189	29	174
Cowley	346	150	300	132	304	134
Crawford	621	249	403	168	337	144
Decatur	52	133	62	174	27	71
Dickinson	261	172	187	126	179	121
Doniphan	96	185	59	114	60	112
Douglas	367	114	284	76	289	68
Edwards	45	148	32	124	38	168
Elk	73	217	46	153	44	163
Ellis	206	126	164	101	145	83
Ellsworth	81	138	82	150	69	132
Finney	190	195	112	108	93	90
Ford	226	170	175	136	136	103
Franklin	212	144	172	121	130	88
Geary	128	154	95	114	109	126
Gove	44	135	28	95	31	106

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2b. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	57	210	43	163	38	159
Grant	42	159	51	179	38	138
Gray	34	104	43	134	24	71
Greeley	18	175	7	59	10	76
Greenwood	161	226	101	155	78	137
Hamilton	19	110	29	172	31	213
Harper	103	168	87	158	45	83
Harvey	286	126	241	95	241	90
Haskell	41	254	37	215	28	155
Hodgeman	13	82	16	87	13	80
Jackson	115	152	86	110	69	85
Jefferson	130	130	123	113	122	103
Jewell	58	156	39	111	26	78
Johnson	1,947	103	1,736	77	1,663	63
Kearny	45	261	25	124	22	109
Kingman	117	157	121	166	73	104
Kiowa	35	131	19	84	11	48
Labette	290	188	211	142	137	94
Lane	24	133	22	157	22	162
Leavenworth	449	178	412	143	425	128
Lincoln	106	288	59	173	38	116
Linn	127	185	90	131	89	131
Logan	41	169	23	89	20	74
Lyon	231	136	169	102	128	77
McPherson	334	143	279	114	222	93
Marion	127	104	116	94	71	65
Marshall	156	148	108	115	110	109
Meade	50	144	39	119	33	86
Miami	287	205	162	103	221	125
Mitchell	91	135	70	108	62	106
Montgomery	414	165	376	153	279	114
Morris	68	141	45	80	55	99
Morton	42	225	40	179	19	85
Nemaha	114	114	94	102	74	72
Neosho	191	159	183	150	119	103
Ness	50	137	35	116	27	81
Norton	48	94	51	100	28	58
Osage	160	145	130	120	90	81

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.
Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2b. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	135	244	76	152	71	147
Ottawa	102	214	37	85	35	87
Pawnee	74	137	56	108	40	77
Phillips	76	128	46	96	26	51
Pottawatomie	154	153	145	137	101	89
Pratt	82	108	97	119	64	84
Rawlins	46	135	46	160	26	79
Reno	518	125	410	95	366	84
Republic	82	147	50	82	51	93
Rice	103	137	94	124	91	133
Riley	250	127	195	90	130	58
Rooks	73	147	40	88	37	81
Rush	83	221	54	146	29	80
Russell	122	167	119	174	80	123
Saline	466	164	386	127	281	88
Scott	44	129	29	83	50	139
Sedgwick	2,631	143	2,194	111	1,990	91
Seward	126	170	75	107	74	113
Shawnee	1,392	164	1,067	119	792	82
Sheridan	35	155	29	130	25	116
Sherman	90	205	61	136	41	97
Smith	62	117	45	91	39	91
Stafford	85	198	51	137	35	86
Stanton	14	127	9	67	7	53
Stevens	49	177	39	123	28	83
Sumner	286	173	232	149	183	116
Thomas	72	140	56	114	43	76
Trego	42	132	44	131	29	93
Wabaunsee	75	186	50	119	36	87
Wallace	24	196	13	74	13	91
Washington	97	145	78	121	50	86
Wichita	17	104	15	79	12	75
Wilson	168	209	162	205	118	157
Woodson	52	169	35	111	29	98
Wyandotte	774	155	589	129	483	107

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2c. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	826	166	737	138	648	105
Frontier	-	-	-	-	-	-
Rural	15	222	13	215	8	100
Dense Rural	55	177	51	162	44	126
Semi-Urban	83	168	75	153	71	134
Urban	672	165	593	134	521	101
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	9	220	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	6	320	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	8	222	-	-	9	212
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	18	193	12	113	13	110
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	25	212	19	152	28	195
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2c. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	36	158	33	68	34	60
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	13	281	7	134	-	-
Lane	-	-	-	-	-	-
Leavenworth	37	199	43	215	33	156
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	9	322	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	21	186	20	209	13	137
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2c. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	11	186	-	-	11	185
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	7	175	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	8	114	15	222	7	96
Scott	-	-	-	-	-	-
Sedgwick	207	161	192	136	179	103
Seward	-	-	-	-	-	-
Shawnee	127	231	112	187	77	117
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	247	146	201	118	185	101

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2d. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	111	187	118	159	120	133
Frontier	-	-	8	334	6	187
Rural	13	340	16	239	14	196
Dense Rural	37	278	32	205	29	147
Semi-Urban	15	133	17	146	9	54
Urban	42	142	44	115	62	134
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	7	329	7	235	7	203
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	6	275	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	6	408	-	-	8	378
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	8	196	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2d. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	8	247	6	141	7	139
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	-	-	6	108
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2d. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	21	173	15	117	33	190
Seward	-	-	-	-	-	-
Shawnee	12	180	14	170	13	134
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2e. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	47	50	69	55	74	43
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	-	-	7	73	6	30
Semi-Urban	12	97	9	64	8	37
Urban	30	46	51	52	58	44
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	7	152	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2e. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	10	41	18	55	16	33
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2e. Age-Adjusted Coronary Heart Disease Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	12	42	20	47	29	52
Seward	-	-	-	-	-	-
Shawnee	-	-	-	-	6	74
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	6	110	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2f. Age-Adjusted Coronary Heart Disease Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	257	97	237	64	269	53
Frontier	13	166	20	163	16	97
Rural	19	110	17	73	18	64
Dense Rural	63	106	46	57	76	66
Semi-Urban	25	72	28	64	28	48
Urban	137	95	126	59	130	44
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	6	132
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	6	106
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	12	88	11	45	18	54
Ford	9	72	-	-	18	95
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2f. Age-Adjusted Coronary Heart Disease Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	7	230	-	-	7	182
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	6	115	6	75	6	63
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	13	56	11	27	16	23
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	6	309	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G2f. Age-Adjusted Coronary Heart Disease Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	6	68	-	-	6	45
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	7	115	7	72
Scott	-	-	-	-	-	-
Sedgwick	45	94	48	62	61	58
Seward	11	131	-	-	9	66
Shawnee	40	143	27	75	26	60
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	34	93	30	60	21	32

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3a. Age-Adjusted Cancer Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	26,591	189	26,691	180	26,842	168
Frontier	1,830	184	1,665	175	1,519	163
Rural	3,156	175	3,140	176	2,943	166
Dense Rural	5,052	189	5,034	185	4,924	175
Semi-Urban	4,267	188	4,372	184	4,398	174
Urban	12,286	194	12,471	178	13,056	165
Allen	206	218	179	196	178	195
Anderson	115	188	113	186	110	175
Atchison	211	205	222	217	183	181
Barber	92	218	72	178	73	197
Barton	345	188	334	177	338	180
Bourbon	196	189	177	176	171	166
Brown	130	165	158	215	129	184
Butler	582	192	607	184	631	175
Chase	55	260	42	213	37	171
Chautauqua	90	231	59	157	66	200
Cherokee	278	211	305	229	282	212
Cheyenne	49	173	38	147	41	135
Clark	25	136	24	127	30	151
Clay	106	154	125	171	121	170
Cloud	147	169	135	166	134	182
Coffey	121	214	131	233	108	179
Comanche	45	217	40	264	33	195
Cowley	470	212	431	195	428	187
Crawford	457	204	455	207	400	179
Decatur	52	169	42	134	65	213
Dickinson	262	196	263	190	261	182
Doniphan	93	185	89	176	82	160
Douglas	608	180	636	163	687	152
Edwards	45	166	49	209	46	213
Elk	78	267	50	195	59	225
Ellis	246	165	257	168	237	146
Ellsworth	83	166	65	132	75	147
Finney	195	158	195	144	215	150
Ford	267	194	214	155	228	151
Franklin	259	186	286	200	261	175
Geary	230	218	232	217	240	211
Gove	35	134	29	122	33	135

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3a. Age-Adjusted Cancer Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	43	162	44	179	29	133
Grant	55	175	56	170	67	195
Gray	52	169	65	207	47	151
Greeley	15	144	14	149	16	147
Greenwood	122	200	137	233	88	158
Hamilton	37	217	26	160	23	155
Harper	112	204	103	199	92	192
Harvey	398	187	380	169	397	164
Haskell	24	133	33	170	25	121
Hodgeman	19	145	22	158	21	134
Jackson	146	198	135	170	154	179
Jefferson	186	185	212	191	212	175
Jewell	57	162	48	148	50	157
Johnson	3,341	169	3,688	158	3,958	146
Kearny	29	151	36	175	46	207
Kingman	122	200	113	185	81	131
Kiowa	31	122	46	215	28	143
Labette	292	197	321	224	288	205
Lane	33	214	29	194	19	127
Leavenworth	580	199	593	183	627	165
Lincoln	51	167	45	157	44	171
Linn	133	198	122	176	157	221
Logan	34	159	38	178	29	132
Lyon	283	169	308	181	322	182
McPherson	318	163	338	163	350	161
Marion	169	160	160	160	152	146
Marshall	159	178	146	172	126	152
Meade	57	180	48	148	54	169
Miami	248	176	279	170	284	155
Mitchell	89	165	92	168	78	151
Montgomery	543	219	488	206	494	209
Morris	95	196	85	179	97	192
Morton	29	152	34	171	34	166
Nemaha	127	149	135	159	126	152
Neosho	227	202	212	188	212	186
Ness	52	164	58	199	38	135
Norton	70	162	77	192	47	110
Osage	190	182	217	208	197	181

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3a. Age-Adjusted Cancer Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	84	221	69	185	60	164
Ottawa	91	227	77	185	62	150
Pawnee	95	188	87	170	88	181
Phillips	82	174	80	170	78	168
Pottawatomie	188	191	184	175	178	156
Pratt	109	168	119	165	125	180
Rawlins	50	170	42	174	40	175
Reno	728	181	746	176	760	173
Republic	87	159	67	117	76	151
Rice	134	186	167	243	121	177
Riley	324	166	317	151	347	152
Rooks	78	181	76	183	64	164
Rush	58	192	61	200	39	117
Russell	111	171	98	163	98	154
Saline	498	170	582	188	584	175
Scott	64	187	48	160	53	151
Sedgwick	4,176	201	4,192	185	4,291	170
Seward	132	158	142	167	142	164
Shawnee	1,846	199	1,796	181	1,866	172
Sheridan	37	174	30	139	31	142
Sherman	78	184	81	178	66	152
Smith	83	188	62	157	63	159
Stafford	64	172	66	194	45	126
Stanton	23	185	24	189	18	127
Stevens	57	198	54	166	48	146
Sumner	321	209	299	195	266	173
Thomas	73	155	73	155	72	147
Trego	50	167	40	156	44	180
Wabaunsee	67	166	84	199	68	149
Wallace	19	166	12	99	18	180
Washington	88	152	76	152	88	168
Wichita	31	196	32	216	29	184
Wilson	133	183	128	175	117	171
Woodson	56	174	68	223	77	289
Wyandotte	1,735	241	1,566	224	1,627	221

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3b. Age-Adjusted Cancer Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	24,590	188	24,342	179	24,083	167
Frontier	1,796	184	1,629	176	1,471	163
Rural	3,111	176	3,065	177	2,841	165
Dense Rural	4,820	191	4,759	186	4,608	177
Semi-Urban	4,062	189	4,081	182	4,079	173
Urban	10,801	192	10,802	176	11,083	163
Allen	198	217	172	195	173	197
Anderson	115	190	111	185	105	168
Atchison	201	207	210	217	175	184
Barber	90	216	71	178	72	197
Barton	333	188	317	174	327	184
Bourbon	187	187	173	178	165	166
Brown	121	161	147	215	113	174
Butler	567	192	581	181	598	172
Chase	54	258	41	214	35	167
Chautauqua	87	229	57	155	62	198
Cherokee	265	209	295	230	271	215
Cheyenne	49	175	38	150	40	133
Clark	25	137	24	130	30	158
Clay	106	155	124	171	120	170
Cloud	146	169	134	167	133	183
Coffey	117	208	129	233	107	181
Comanche	44	211	39	263	33	198
Cowley	438	207	397	191	385	181
Crawford	448	204	448	209	391	181
Decatur	52	170	42	136	65	215
Dickinson	253	194	253	186	253	181
Doniphan	91	187	86	175	77	157
Douglas	570	181	594	163	647	156
Edwards	44	169	49	226	43	226
Elk	76	264	48	190	58	226
Ellis	245	166	253	167	232	146
Ellsworth	81	168	64	135	72	147
Finney	162	165	152	150	161	160
Ford	249	210	195	170	200	167
Franklin	255	188	265	190	250	174
Geary	183	221	181	218	182	216
Gove	35	135	29	124	33	137

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3b. Age-Adjusted Cancer Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	38	151	41	176	25	114
Grant	51	187	51	182	54	197
Gray	52	175	62	206	45	152
Greeley	15	146	14	154	15	146
Greenwood	122	202	135	234	84	155
Hamilton	35	214	25	161	23	162
Harper	111	204	103	202	88	185
Harvey	386	192	362	172	377	165
Haskell	24	144	31	177	23	124
Hodgeman	18	140	22	162	21	138
Jackson	139	200	127	170	133	165
Jefferson	184	186	205	188	208	175
Jewell	57	163	48	149	49	153
Johnson	3,216	171	3,491	160	3,685	146
Kearny	27	155	33	181	45	239
Kingman	121	200	113	188	81	133
Kiowa	30	120	45	214	28	146
Labette	274	197	303	226	272	208
Lane	33	215	29	197	18	122
Leavenworth	526	201	547	186	551	162
Lincoln	51	169	45	159	44	175
Linn	133	200	120	176	154	222
Logan	34	162	38	182	29	135
Lyon	266	171	286	184	306	193
McPherson	315	164	332	163	343	162
Marion	168	161	155	157	144	140
Marshall	158	177	144	171	125	150
Meade	54	176	47	148	52	170
Miami	238	173	264	166	275	155
Mitchell	88	163	92	169	75	147
Montgomery	499	219	434	197	453	210
Morris	95	199	80	171	95	193
Morton	29	160	30	162	29	150
Nemaha	126	149	134	160	126	154
Neosho	221	202	207	189	205	186
Ness	52	164	58	200	38	137
Norton	70	167	75	192	46	109
Osage	189	183	212	206	189	178

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3b. Age-Adjusted Cancer Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	84	222	68	183	60	166
Ottawa	90	227	76	185	60	148
Pawnee	92	191	83	172	82	184
Phillips	82	175	80	173	77	169
Pottawatomie	186	193	180	175	169	152
Pratt	107	168	113	160	116	169
Rawlins	50	171	41	173	39	176
Reno	708	185	702	174	694	167
Republic	87	160	66	117	75	152
Rice	130	186	160	244	118	183
Riley	306	167	292	149	319	151
Rooks	78	183	75	182	64	166
Rush	58	193	61	203	38	115
Russell	111	173	96	162	98	157
Saline	472	170	552	190	540	174
Scott	61	180	47	162	50	147
Sedgwick	3,656	200	3,537	181	3,546	167
Seward	112	169	121	190	108	172
Shawnee	1,647	200	1,577	180	1,626	172
Sheridan	37	175	30	140	30	139
Sherman	75	181	77	176	61	145
Smith	83	188	62	158	63	160
Stafford	64	175	65	196	43	125
Stanton	19	165	22	194	14	108
Stevens	55	211	53	175	44	146
Sumner	312	210	283	191	256	173
Thomas	73	156	73	157	72	150
Trego	50	169	40	157	44	182
Wabaunsee	65	164	83	201	67	150
Wallace	18	164	12	101	17	163
Washington	88	153	76	153	88	169
Wichita	29	203	28	201	27	199
Wilson	131	184	123	172	112	168
Woodson	56	177	67	223	76	291
Wyandotte	1,186	244	1,056	236	1,028	231

Note: Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3c. Age-Adjusted Cancer Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	1,375	266	1,311	233	1,413	218
Frontier	10	430	6	236	6	214
Rural	6	94	15	234	9	127
Dense Rural	84	271	67	220	63	178
Semi-Urban	128	250	147	272	121	200
Urban	1,147	269	1,074	230	1,213	224
Allen	6	436	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	7	157	7	174	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	6	234	-	-	-	-
Brown	-	-	-	-	-	-
Butler	7	373	9	358	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	14	341	10	260	10	244
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	30	298	15	143	19	146
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	35	270	36	238	37	199
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3c. Age-Adjusted Cancer Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	56	216	75	195	111	189
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	16	339	12	268	9	175
Lane	-	-	-	-	-	-
Leavenworth	49	241	33	173	51	231
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	6	280	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	6	326	11	447	-	-
Mitchell	-	-	-	-	-	-
Montgomery	35	328	39	429	29	310
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3c. Age-Adjusted Cancer Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	10	161	13	228	16	239
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	7	139	8	193	10	130
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	20	313	18	264	12	152
Scott	-	-	-	-	-	-
Sedgwick	380	277	386	251	425	235
Seward	7	216	6	297	8	237
Shawnee	135	243	143	236	147	229
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	497	284	422	234	460	233

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3d. Age-Adjusted Cancer Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	146	211	182	229	205	217
Frontier	8	398	8	248	6	189
Rural	15	297	17	236	22	242
Dense Rural	38	212	51	306	57	245
Semi-Urban	13	120	24	175	26	163
Urban	72	222	82	211	94	222
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	6	2,940	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	8	287	7	219	12	319
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	8	314	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	9	277	9	385	19	758
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	12	449	8	113
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	7	915	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3d. Age-Adjusted Cancer Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	6	176	8	208	15	296
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	8	264	11	230	12	225
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3d. Age-Adjusted Cancer Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	8	464
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	30	194	38	196	45	267
Seward	-	-	-	-	-	-
Shawnee	20	357	9	120	16	194
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	7	135	10	270	9	206

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3e. Age-Adjusted Cancer Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	121	93	171	103	209	97
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	13	85	17	151	23	131
Semi-Urban	23	132	20	82	36	171
Urban	82	85	133	104	147	85
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	7	247
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	9	154	8	110	14	219
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3e. Age-Adjusted Cancer Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	26	83	38	78	49	72
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	8	157
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3e. Age-Adjusted Cancer Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	6	207	-	-	8	209
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	36	94	71	134	58	80
Seward	-	-	-	-	-	-
Shawnee	6	72	-	-	7	77
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	12	161	11	126	21	189

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3f. Age-Adjusted Cancer Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	355	113	479	111	650	117
Frontier	15	166	17	121	23	127
Rural	21	117	28	111	47	147
Dense Rural	97	128	108	100	133	102
Semi-Urban	40	102	63	130	72	112
Urban	182	108	263	110	375	120
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	8	121	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	6	165	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	6	156	8	157	7	95
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	28	123	34	134	43	120
Ford	14	88	12	66	21	108
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3f. Age-Adjusted Cancer Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	12	210
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	8	99	11	118	12	100
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	35	115	46	94	64	107
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	10	138
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	9	105	12	68	8	56
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	6	162	7	151
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G3f. Age-Adjusted Cancer Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	6	60	19	169	19	128
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	6	193	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	7	100	18	210
Scott	-	-	-	-	-	-
Sedgwick	73	130	99	119	141	124
Seward	11	138	13	84	22	129
Shawnee	38	126	50	126	61	136
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	7	186	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	32	72	61	114	94	127

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4a. Age-Adjusted Diabetes Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	3,463	24	3,546	23	3,273	20
Frontier	245	23	263	26	205	21
Rural	475	24	532	28	440	24
Dense Rural	728	26	729	26	678	24
Semi-Urban	511	22	559	23	544	21
Urban	1,504	24	1,462	21	1,406	18
Allen	32	31	34	35	29	28
Anderson	13	18	22	33	9	14
Atchison	38	36	39	37	36	35
Barber	8	20	9	25	10	25
Barton	47	24	33	17	41	22
Bourbon	16	15	21	20	26	25
Brown	15	21	25	30	32	46
Butler	76	25	78	24	62	17
Chase	-	-	-	-	-	-
Chautauqua	-	-	9	21	7	19
Cherokee	32	25	43	32	50	37
Cheyenne	-	-	-	-	6	33
Clark	-	-	-	-	-	-
Clay	18	20	15	18	13	18
Cloud	20	20	36	40	13	16
Coffey	16	28	20	34	12	22
Comanche	-	-	-	-	-	-
Cowley	100	44	92	41	84	36
Crawford	60	26	55	24	59	27
Decatur	15	44	14	35	-	-
Dickinson	23	17	22	15	23	14
Doniphan	12	24	13	28	9	19
Douglas	84	25	82	21	85	19
Edwards	6	22	8	29	9	36
Elk	-	-	6	20	6	21
Ellis	42	27	44	28	32	19
Ellsworth	11	20	22	38	31	54
Finney	36	30	34	26	33	24
Ford	28	20	36	26	34	23
Franklin	31	22	43	29	37	26
Geary	35	34	27	26	36	32
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4a. Age-Adjusted Diabetes Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	9	32	6	23
Grant	-	-	11	34	-	-
Gray	7	22	7	22	6	19
Greeley	9	82	-	-	-	-
Greenwood	15	24	13	22	8	13
Hamilton	-	-	-	-	-	-
Harper	13	21	17	32	28	56
Harvey	38	17	63	26	59	22
Haskell	6	33	10	52	6	30
Hodgeman	-	-	-	-	-	-
Jackson	15	20	15	20	10	12
Jefferson	27	27	16	15	18	15
Jewell	8	23	10	29	-	-
Johnson	302	15	313	13	326	12
Kearny	9	47	7	33	13	63
Kingman	31	43	27	36	21	33
Kiowa	12	46	-	-	-	-
Labette	33	22	33	22	32	22
Lane	-	-	-	-	-	-
Leavenworth	74	26	86	26	100	26
Lincoln	-	-	7	23	-	-
Linn	12	17	9	12	8	12
Logan	6	22	7	27	-	-
Lyon	47	27	32	16	36	20
McPherson	38	18	35	16	39	17
Marion	16	12	33	29	20	17
Marshall	8	10	10	11	9	9
Meade	19	59	13	39	12	37
Miami	20	14	21	13	18	10
Mitchell	10	15	16	25	13	27
Montgomery	73	28	69	28	60	25
Morris	14	28	8	14	7	11
Morton	-	-	-	-	-	-
Nemaha	26	27	25	30	20	22
Neosho	27	24	20	18	11	10
Ness	7	16	15	43	9	22
Norton	12	31	9	18	-	-
Osage	19	18	27	25	20	19

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4a. Age-Adjusted Diabetes Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	18	40	31	66	8	22
Ottawa	9	23	13	27	10	24
Pawnee	31	50	20	38	6	11
Phillips	16	30	11	22	12	27
Pottawatomie	20	20	24	22	23	21
Pratt	15	22	21	28	21	32
Rawlins	12	37	8	28	-	-
Reno	89	21	88	20	106	23
Republic	16	27	8	18	14	20
Rice	19	25	22	28	19	27
Riley	40	21	35	16	39	16
Rooks	12	27	15	32	8	17
Rush	-	-	-	-	8	20
Russell	24	35	18	28	17	30
Saline	49	17	80	26	68	20
Scott	8	19	-	-	-	-
Sedgwick	513	25	548	24	488	19
Seward	30	36	42	50	30	35
Shawnee	262	28	194	20	207	20
Sheridan	-	-	-	-	-	-
Sherman	-	-	20	51	11	28
Smith	-	-	-	-	-	-
Stafford	9	21	8	26	13	35
Stanton	-	-	-	-	-	-
Stevens	10	33	12	34	11	36
Sumner	66	40	74	48	62	39
Thomas	10	21	9	17	10	21
Trego	-	-	-	-	-	-
Wabaunsee	15	36	18	43	12	27
Wallace	-	-	-	-	-	-
Washington	9	16	8	13	-	-
Wichita	6	36	-	-	-	-
Wilson	35	45	37	49	35	50
Woodson	-	-	8	26	6	21
Wyandotte	269	37	239	34	200	28

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4b. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	2,990	22	3,011	22	2,783	19
Frontier	237	22	254	26	191	20
Rural	461	24	507	27	411	23
Dense Rural	660	25	655	25	614	23
Semi-Urban	458	21	493	21	481	20
Urban	1,174	21	1,102	18	1,086	16
Allen	29	28	31	32	28	29
Anderson	13	19	22	34	8	12
Atchison	33	32	30	30	32	33
Barber	8	20	9	26	9	22
Barton	45	23	32	17	35	19
Bourbon	15	15	21	21	26	26
Brown	12	18	21	27	24	34
Butler	72	24	74	23	57	16
Chase	-	-	-	-	-	-
Chautauqua	-	-	9	22	-	-
Cherokee	30	25	42	32	47	36
Cheyenne	-	-	-	-	6	34
Clark	-	-	-	-	-	-
Clay	18	20	15	18	13	18
Cloud	20	20	36	40	12	15
Coffey	16	29	20	35	12	22
Comanche	-	-	-	-	-	-
Cowley	87	40	83	39	68	31
Crawford	58	26	54	24	54	26
Decatur	15	44	14	35	-	-
Dickinson	22	17	21	15	21	13
Doniphan	10	19	12	26	9	20
Douglas	71	22	71	20	76	18
Edwards	-	-	8	31	6	24
Elk	-	-	6	20	6	21
Ellis	42	28	44	28	31	19
Ellsworth	11	20	22	39	30	53
Finney	23	23	21	20	22	21
Ford	21	17	29	25	28	27
Franklin	30	22	41	29	34	25
Geary	26	31	15	19	25	30
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4b. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	8	30	6	24
Grant	-	-	7	25	-	-
Gray	7	23	6	19	6	20
Greeley	9	83	-	-	-	-
Greenwood	15	24	13	23	8	13
Hamilton	-	-	-	-	-	-
Harper	13	21	16	31	27	55
Harvey	36	16	58	25	52	21
Haskell	6	37	7	40	-	-
Hodgeman	-	-	-	-	-	-
Jackson	14	20	10	14	9	11
Jefferson	26	26	15	14	18	15
Jewell	8	23	10	29	-	-
Johnson	289	15	276	13	298	12
Kearny	7	39	7	36	12	65
Kingman	31	44	27	37	21	33
Kiowa	12	46	-	-	-	-
Labette	28	19	32	22	31	23
Lane	-	-	-	-	-	-
Leavenworth	59	23	67	23	84	25
Lincoln	-	-	7	23	-	-
Linn	12	17	9	13	8	12
Logan	6	23	-	-	-	-
Lyon	42	26	29	15	32	19
McPherson	38	18	34	16	38	17
Marion	16	13	33	29	17	14
Marshall	8	10	10	11	9	9
Meade	17	55	13	40	12	39
Miami	19	14	18	11	16	10
Mitchell	10	15	16	26	12	22
Montgomery	61	26	59	27	52	24
Morris	14	28	8	14	7	11
Morton	-	-	-	-	-	-
Nemaha	25	26	24	29	20	23
Neosho	25	23	18	16	10	9
Ness	7	16	15	44	9	22
Norton	12	32	9	18	-	-
Osage	19	18	27	26	20	19

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4b. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	18	40	31	67	8	22
Ottawa	9	23	12	25	9	21
Pawnee	29	48	18	36	6	12
Phillips	16	30	10	20	11	25
Pottawatomie	20	20	23	22	23	22
Pratt	15	23	20	27	20	32
Rawlins	12	37	8	28	-	-
Reno	76	19	78	19	94	22
Republic	16	27	8	18	14	20
Rice	17	23	19	25	18	26
Riley	36	20	29	14	36	15
Rooks	12	28	15	33	8	18
Rush	-	-	-	-	8	20
Russell	23	33	17	27	17	30
Saline	44	16	67	23	61	20
Scott	8	19	-	-	-	-
Sedgwick	405	22	415	21	362	17
Seward	29	41	32	51	26	42
Shawnee	207	25	150	17	159	17
Sheridan	-	-	-	-	-	-
Sherman	-	-	18	48	11	31
Smith	-	-	-	-	-	-
Stafford	9	21	8	27	13	36
Stanton	-	-	-	-	-	-
Stevens	8	28	11	32	10	34
Sumner	62	38	69	46	60	39
Thomas	10	21	9	17	9	19
Trego	-	-	-	-	-	-
Wabaunsee	15	37	18	44	12	28
Wallace	-	-	-	-	-	-
Washington	9	16	8	13	-	-
Wichita	-	-	-	-	-	-
Wilson	32	42	37	50	32	47
Woodson	-	-	8	27	6	21
Wyandotte	143	30	123	27	107	25

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4c. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	314	62	306	55	246	39
Frontier	-	-	-	-	-	-
Rural	6	93	9	135	-	-
Dense Rural	27	85	20	64	18	50
Semi-Urban	29	59	40	83	30	55
Urban	251	61	236	50	193	36
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	8	192	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	10	198
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	9	93	6	66	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	8	72	10	81	9	62
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4c. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	7	31	18	42	10	21
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	14	73	12	50	10	47
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	7	63	6	63	6	66
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4c. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	6	105	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	74	55	79	50	71	39
Seward	-	-	-	-	-	-
Shawnee	41	74	27	44	28	42
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	106	62	94	53	70	39

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4d. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	26	35	39	48	44	44
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	9	92
Dense Rural	6	32	9	47	11	51
Semi-Urban	-	-	6	63	-	-
Urban	14	42	20	46	16	31
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	8	204
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4d. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	-	-	-	-
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4d. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	-	-	7	35	7	45
Seward	-	-	-	-	-	-
Shawnee	-	-	-	-	-	-
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4e. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	15	18	26	22	20	10
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	-	-	7	88	-	-
Semi-Urban	-	-	-	-	-	-
Urban	12	20	17	17	15	9
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4e. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	-	-	-	-
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4e. Age-Adjusted Diabetes Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	10	42	11	27	8	10
Seward	-	-	-	-	-	-
Shawnee	-	-	-	-	-	-
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4f. Age-Adjusted Diabetes Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	118	41	138	35	130	25
Frontier	7	89	6	37	9	47
Rural	-	-	12	47	9	33
Dense Rural	35	55	32	37	29	26
Semi-Urban	19	54	17	33	17	30
Urban	53	30	70	32	66	22
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	13	100	10	46	8	26
Ford	-	-	6	27	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4f. Age-Adjusted Diabetes Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	7	19	9	13
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G4f. Age-Adjusted Diabetes Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	6	62	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	7	91	-	-
Scott	-	-	-	-	-	-
Sedgwick	20	31	27	35	28	25
Seward	-	-	6	48	-	-
Shawnee	9	27	12	35	13	29
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	19	46	19	38	12	18

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5a. Age-Adjusted Unintentional Injury Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	5,457	39	5,863	41	6,544	43
Frontier	416	60	436	65	468	69
Rural	769	53	752	54	809	59
Dense Rural	1,193	47	1,205	48	1,343	53
Semi-Urban	905	40	989	43	1,131	47
Urban	2,171	32	2,476	34	2,792	35
Allen	44	52	47	66	45	64
Anderson	22	57	21	52	19	47
Atchison	40	44	32	35	55	58
Barber	15	54	22	76	20	82
Barton	68	41	59	39	74	49
Bourbon	53	62	49	66	44	54
Brown	34	53	45	78	28	52
Butler	132	44	162	50	161	48
Chase	7	55	10	62	7	43
Chautauqua	14	54	18	79	17	64
Cherokee	59	50	69	63	59	55
Cheyenne	12	80	14	68	14	50
Clark	6	42	10	97	-	-
Clay	23	36	35	70	29	55
Cloud	32	45	25	44	36	59
Coffey	20	38	26	50	39	91
Comanche	6	49	7	96	13	108
Cowley	103	52	96	50	120	63
Crawford	89	42	98	46	102	49
Decatur	9	40	11	68	16	86
Dickinson	46	44	45	41	60	53
Doniphan	23	50	21	43	14	31
Douglas	126	29	155	35	169	34
Edwards	16	104	8	38	18	111
Elk	18	84	13	77	10	67
Ellis	49	33	46	31	73	47
Ellsworth	26	63	15	39	19	52
Finney	80	43	86	55	87	53
Ford	65	41	55	34	89	54
Franklin	55	42	62	48	68	51
Geary	55	42	56	42	72	48
Gove	10	61	14	71	13	86

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5a. Age-Adjusted Unintentional Injury Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	15	106	14	97	7	54
Grant	14	39	27	75	8	22
Gray	18	58	21	70	17	54
Greeley	6	85	-	-	8	119
Greenwood	29	66	24	58	22	62
Hamilton	7	51	10	70	15	110
Harper	30	71	30	90	25	71
Harvey	78	42	83	42	115	50
Haskell	12	62	13	67	12	61
Hodgeman	9	64	-	-	7	48
Jackson	30	45	47	69	36	47
Jefferson	50	54	62	64	53	56
Jewell	10	49	8	38	9	45
Johnson	456	21	560	23	634	23
Kearny	17	81	16	81	18	88
Kingman	27	58	24	47	38	75
Kiowa	9	53	19	98	12	76
Labette	65	53	72	58	76	68
Lane	8	59	7	69	8	64
Leavenworth	113	34	115	33	133	34
Lincoln	18	108	6	38	10	43
Linn	39	83	24	50	27	51
Logan	10	62	8	55	9	59
Lyon	72	40	78	43	74	42
McPherson	69	36	74	38	97	48
Marion	55	71	46	60	42	47
Marshall	33	50	25	39	25	40
Meade	14	57	15	62	12	54
Miami	73	51	64	41	58	36
Mitchell	12	28	18	42	20	49
Montgomery	136	70	128	69	107	55
Morris	19	49	19	64	16	45
Morton	6	35	10	61	14	81
Nemaha	34	50	32	48	37	52
Neosho	54	60	52	53	44	50
Ness	11	56	15	74	18	108
Norton	9	26	14	38	17	48
Osage	51	61	51	59	57	66

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5a. Age-Adjusted Unintentional Injury Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	10	41	15	59	10	44
Ottawa	13	42	14	36	14	47
Pawnee	22	57	19	46	25	66
Phillips	14	44	13	38	19	63
Pottawatomie	49	52	46	45	48	43
Pratt	20	35	22	34	36	61
Rawlins	8	39	15	105	-	-
Reno	120	34	168	47	234	58
Republic	25	57	12	51	19	61
Rice	32	53	40	70	31	53
Riley	69	29	67	24	86	31
Rooks	18	66	25	85	19	69
Rush	11	46	13	70	17	74
Russell	21	46	21	49	30	66
Saline	98	35	101	34	128	42
Scott	12	34	21	81	27	95
Sedgwick	851	39	1,016	44	1,184	48
Seward	44	41	50	47	55	55
Shawnee	323	37	339	37	385	40
Sheridan	8	43	7	51	11	93
Sherman	16	46	18	54	24	64
Smith	15	52	11	37	23	93
Stafford	11	43	15	72	21	80
Stanton	16	134	9	84	8	62
Stevens	12	45	26	93	19	70
Sumner	79	59	68	54	83	63
Thomas	17	41	23	52	29	62
Trego	8	41	12	51	13	57
Wabaunsee	17	47	19	52	25	66
Wallace	8	74	-	-	7	65
Washington	28	76	15	41	26	74
Wichita	-	-	6	56	9	75
Wilson	53	88	32	56	31	65
Woodson	24	88	15	77	24	144
Wyandotte	302	40	291	39	287	38

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5b. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	4,827	40	5,073	41	5,648	44
Frontier	390	58	412	66	429	67
Rural	747	54	698	53	755	59
Dense Rural	1,058	47	1,046	48	1,174	54
Semi-Urban	844	42	894	43	1,011	46
Urban	1,785	31	2,020	34	2,278	36
Allen	43	53	45	66	44	66
Anderson	22	58	21	54	19	48
Atchison	38	46	28	32	52	59
Barber	14	52	21	72	20	86
Barton	59	38	57	42	68	51
Bourbon	52	64	44	62	43	56
Brown	32	55	39	72	23	47
Butler	129	45	150	49	148	47
Chase	7	58	10	66	6	36
Chautauqua	14	57	18	85	17	68
Cherokee	58	52	61	59	57	58
Cheyenne	11	73	13	59	14	51
Clark	6	44	9	94	-	-
Clay	21	33	34	69	29	57
Cloud	32	46	25	45	35	59
Coffey	19	36	25	50	37	92
Comanche	6	50	6	84	13	112
Cowley	92	51	88	52	106	64
Crawford	85	42	96	48	88	45
Decatur	9	40	10	60	16	89
Dickinson	46	46	42	40	55	50
Doniphan	23	53	21	45	14	34
Douglas	107	28	136	34	157	36
Edwards	15	115	8	43	13	92
Elk	17	80	13	81	10	70
Ellis	48	33	46	32	70	47
Ellsworth	25	66	14	39	19	59
Finney	33	32	48	49	58	62
Ford	52	47	38	41	53	54
Franklin	53	43	54	43	61	47
Geary	39	44	37	41	48	47
Gove	10	62	14	73	12	78

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5b. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	13	89	12	80	-	-
Grant	11	43	21	93	-	-
Gray	17	59	16	59	17	59
Greeley	-	-	-	-	7	116
Greenwood	29	68	24	60	22	65
Hamilton	6	51	7	48	13	126
Harper	30	73	29	93	21	61
Harvey	73	43	80	44	108	51
Haskell	11	72	12	81	10	70
Hodgeman	9	66	-	-	7	51
Jackson	23	35	39	62	30	43
Jefferson	49	54	59	63	52	57
Jewell	10	50	8	38	9	46
Johnson	421	21	490	22	573	24
Kearny	7	46	15	101	13	91
Kingman	26	56	24	49	38	78
Kiowa	7	39	19	100	11	71
Labette	60	54	64	56	72	73
Lane	8	59	7	73	8	68
Leavenworth	97	35	106	36	120	37
Lincoln	18	110	6	40	10	44
Linn	38	83	24	52	26	50
Logan	10	64	8	57	9	63
Lyon	62	42	61	40	64	45
McPherson	68	37	73	39	89	44
Marion	55	74	42	57	41	46
Marshall	33	51	25	39	24	38
Meade	14	63	12	54	7	32
Miami	72	53	62	41	55	36
Mitchell	12	28	17	42	20	50
Montgomery	127	75	110	69	98	61
Morris	19	51	19	67	16	48
Morton	6	40	9	68	11	71
Nemaha	34	51	31	47	34	47
Neosho	51	59	50	53	42	50
Ness	11	58	15	82	17	108
Norton	9	29	12	35	17	52
Osage	50	62	49	59	56	68

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5b. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	10	41	15	60	10	45
Ottawa	13	43	14	36	14	49
Pawnee	19	53	19	51	23	71
Phillips	14	45	13	39	19	66
Pottawatomie	45	49	45	47	45	42
Pratt	19	35	21	34	33	59
Rawlins	8	39	15	109	-	-
Reno	115	36	154	46	215	58
Republic	25	58	11	47	19	63
Rice	28	49	37	70	28	53
Riley	59	29	63	26	74	31
Rooks	17	63	25	88	18	67
Rush	10	39	13	73	17	77
Russell	21	47	21	51	27	63
Saline	92	37	88	32	116	43
Scott	12	34	18	78	26	113
Sedgwick	716	40	818	44	930	48
Seward	31	53	23	45	24	47
Shawnee	269	36	284	37	327	41
Sheridan	8	43	7	53	8	59
Sherman	14	43	18	61	22	62
Smith	15	53	11	37	23	95
Stafford	11	47	13	70	19	75
Stanton	15	155	6	94	6	60
Stevens	12	60	16	69	12	59
Sumner	75	60	65	55	80	65
Thomas	17	42	23	54	25	54
Trego	8	42	11	49	13	59
Wabaunsee	16	46	18	49	23	62
Wallace	8	79	-	-	7	71
Washington	28	77	14	37	25	71
Wichita	-	-	-	-	8	80
Wilson	53	91	29	52	28	63
Woodson	24	90	14	72	23	148
Wyandotte	175	41	186	49	171	46

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5c. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	272	38	284	37	304	37
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	15	36	20	52	18	39
Semi-Urban	29	30	35	39	44	52
Urban	222	38	223	36	236	35
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	8	47	10	55	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	9	24	15	63	16	71
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5c. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	13	20	27	29	26	17
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	14	50	7	15	8	34
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	6	55	10	91	6	62
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5c. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	7	22	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	64	34	84	43	98	46
Seward	-	-	-	-	6	162
Shawnee	36	48	29	40	31	37
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	87	42	66	33	68	34

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5d. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	45	37	62	57	56	46
Frontier	-	-	-	-	-	-
Rural	-	-	6	60	10	102
Dense Rural	11	39	15	69	11	42
Semi-Urban	-	-	13	80	6	24
Urban	22	41	27	46	29	52
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	9	181	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5d. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	6	145	6	138	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	-	-	-	-
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5d. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	9	42	13	61	18	90
Seward	-	-	-	-	-	-
Shawnee	-	-	-	-	6	55
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5e. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	45	20	27	13	48	18
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	12	85	-	-	6	19
Semi-Urban	6	17	6	20	9	34
Urban	25	15	18	11	32	16
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5e. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	6	7	6	8	6	9
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5e. Age-Adjusted Unintentional Injury Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	12	20	11	19	16	23
Seward	-	-	-	-	-	-
Shawnee	-	-	-	-	-	-
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5f. Age-Adjusted Unintentional Injury Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	265	30	314	31	372	32
Frontier	21	57	18	49	32	79
Rural	12	30	29	51	27	41
Dense Rural	97	39	102	40	116	40
Semi-Urban	20	26	23	19	42	36
Urban	115	25	141	27	155	25
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	7	46	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	6	77	6	70	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	43	61	32	50	25	41
Ford	12	24	16	25	35	50
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5f. Age-Adjusted Unintentional Injury Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	6	62	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	13	20	25	20	17	11
Kearny	10	151	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	8	32	13	68	7	22
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G5f. Age-Adjusted Unintentional Injury Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	7	35
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	8	55
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	7	35
Scott	-	-	-	-	-	-
Sedgwick	49	26	63	33	84	41
Seward	13	25	23	32	24	58
Shawnee	15	26	14	30	13	13
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	8	134	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	36	30	37	26	36	25

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6a. Age-Adjusted Homicide Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	666	5	565	4	585	4
Frontier	16	3	15	3	19	4
Rural	18	2	21	2	21	2
Dense Rural	84	4	61	3	60	3
Semi-Urban	57	3	56	2	76	3
Urban	491	7	412	5	407	5
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	7	5	6	5	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	9	3	-	-	6	2
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	6	4	-	-	-	-
Crawford	14	7	-	-	9	5
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	12	3	12	2	6	1
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	9	5	7	4	6	3
Ford	15	9	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	8	5	11	6	13	7
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6a. Age-Adjusted Homicide Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	42	2	52	2	48	2
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	13	4	7	2	18	5
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	10	5	6	3	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	10	6	9	6
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6a. Age-Adjusted Homicide Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	7	2	8	3	8	3
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	6	1	7	2
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	8	3	18	7
Scott	-	-	-	-	-	-
Sedgwick	132	6	152	6	120	5
Seward	-	-	-	-	-	-
Shawnee	91	11	48	6	78	10
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	201	25	141	18	137	18

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6b. Age-Adjusted Homicide Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	312	3	234	2	282	3
Frontier	8	2	14	3	15	4
Rural	18	2	17	2	17	2
Dense Rural	50	3	41	2	40	2
Semi-Urban	46	2	40	2	51	3
Urban	190	3	122	2	159	3
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	7	2	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	14	7	-	-	8	5
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	9	3	8	2	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	9	9	9	8
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6b. Age-Adjusted Homicide Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	34	2	29	1	29	1
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	10	3	-	-	12	4
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	6	5	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6b. Age-Adjusted Homicide Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	7	3	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	6	3	14	6
Scott	-	-	-	-	-	-
Sedgwick	52	3	46	3	49	3
Seward	-	-	-	-	-	-
Shawnee	40	6	16	3	37	6
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	45	11	19	5	27	8

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6c. Age-Adjusted Homicide Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	252	30	223	25	178	19
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	-	-	-	-	6	9
Semi-Urban	9	9	11	9	10	9
Urban	240	36	209	30	160	21
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6c. Age-Adjusted Homicide Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	9	8	14	9
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6c. Age-Adjusted Homicide Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	62	28	74	33	41	17
Seward	-	-	-	-	-	-
Shawnee	40	50	26	31	27	34
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	130	58	97	47	74	39

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6d. Age-Adjusted Homicide Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	-	-	6	4	8	6
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	-	-	-	-	-	-
Semi-Urban	-	-	-	-	-	-
Urban	-	-	-	-	-	-
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6d. Age-Adjusted Homicide Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	-	-	-	-
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6d. Age-Adjusted Homicide Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	-	-	-	-	-	-
Seward	-	-	-	-	-	-
Shawnee	-	-	-	-	-	-
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6e. Age-Adjusted Homicide Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	12	4	12	4	11	3
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	-	-	-	-	-	-
Semi-Urban	-	-	-	-	-	-
Urban	8	3	10	4	8	3
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6e. Age-Adjusted Homicide Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	-	-	-	-
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6e. Age-Adjusted Homicide Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	6	6	-	-	7	6
Seward	-	-	-	-	-	-
Shawnee	-	-	-	-	-	-
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6f. Age-Adjusted Homicide Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	84	8	85	7	87	5
Frontier	7	16	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	26	8	13	4	11	3
Semi-Urban	-	-	-	-	7	5
Urban	49	8	63	9	64	6
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	6	9	-	-	-	-
Ford	13	16	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6f. Age-Adjusted Homicide Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	9	9	-	-
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G6f. Age-Adjusted Homicide Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	12	5	23	9	17	4
Seward	-	-	-	-	-	-
Shawnee	9	13	6	5	8	7
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	24	18	23	12	33	15

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7a. Age-Adjusted Suicide Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	1,616	12	1,832	13	2,100	15
Frontier	88	14	83	14	99	19
Rural	149	12	192	16	184	16
Dense Rural	283	12	293	13	375	16
Semi-Urban	268	13	321	15	316	14
Urban	828	12	942	13	1,125	14
Allen	12	16	9	13	16	24
Anderson	-	-	7	20	7	22
Atchison	10	13	9	12	13	18
Barber	10	40	-	-	7	31
Barton	16	11	13	11	42	34
Bourbon	8	10	10	12	15	23
Brown	-	-	12	24	9	21
Butler	31	11	32	10	53	16
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	17	15	16	15	22	20
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	7	17	11	26	9	23
Cloud	-	-	-	-	7	15
Coffey	6	14	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	30	17	27	15	23	14
Crawford	34	18	38	20	20	10
Decatur	-	-	-	-	-	-
Dickinson	11	12	9	10	12	12
Doniphan	-	-	8	19	-	-
Douglas	44	10	64	13	71	13
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	18	13	26	18	26	19
Ellsworth	-	-	6	19	-	-
Finney	11	7	18	12	20	11
Ford	12	8	15	10	24	15
Franklin	16	13	25	19	18	13
Geary	23	17	25	17	31	17
Gove	-	-	-	-	6	40

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7a. Age-Adjusted Suicide Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	8	18	-	-	8	32
Hamilton	-	-	-	-	-	-
Harper	-	-	7	22	-	-
Harvey	15	9	30	18	22	13
Haskell	6	30	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	12	19
Jefferson	10	11	16	16	12	13
Jewell	-	-	-	-	-	-
Johnson	243	11	269	11	384	14
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	12	11	17	17	12	10
Lane	-	-	-	-	-	-
Leavenworth	46	14	49	13	61	16
Lincoln	6	28	-	-	-	-
Linn	-	-	10	21	11	21
Logan	-	-	-	-	-	-
Lyon	25	14	17	10	30	18
McPherson	15	11	16	11	19	12
Marion	7	9	13	18	9	15
Marshall	-	-	10	19	-	-
Meade	-	-	-	-	-	-
Miami	24	16	27	17	25	15
Mitchell	7	19	7	21	-	-
Montgomery	31	18	24	15	26	15
Morris	-	-	-	-	6	26
Morton	-	-	-	-	-	-
Nemaha	8	14	8	17	9	20
Neosho	17	19	9	12	10	10
Ness	-	-	-	-	-	-
Norton	-	-	-	-	8	27
Osage	9	11	16	20	12	14

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7a. Age-Adjusted Suicide Mortality Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	6	18	-	-
Pawnee	8	20	-	-	8	22
Phillips	-	-	-	-	6	21
Pottawatomie	12	14	13	13	20	17
Pratt	11	24	9	19	11	26
Rawlins	-	-	-	-	-	-
Reno	41	12	47	15	45	14
Republic	-	-	6	19	-	-
Rice	-	-	12	21	8	14
Riley	25	8	37	13	36	11
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	7	22	9	24	9	24
Saline	28	10	36	13	40	15
Scott	-	-	-	-	-	-
Sedgwick	247	11	320	14	344	14
Seward	10	9	6	5	14	12
Shawnee	134	16	130	15	144	16
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	20	17	19	16	19	16
Thomas	6	18	8	19	-	-
Trego	-	-	-	-	-	-
Wabaunsee	7	18	10	31	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	6	12	6	13	7	16
Woodson	-	-	-	-	-	-
Wyandotte	114	15	110	15	121	15

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7b. Age-Adjusted Suicide Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	1,499	13	1,636	14	1,871	16
Frontier	86	14	81	15	91	19
Rural	149	13	182	16	171	16
Dense Rural	263	13	267	14	337	18
Semi-Urban	256	13	296	16	284	14
Urban	745	13	809	14	987	16
Allen	12	17	8	12	15	24
Anderson	-	-	6	17	7	23
Atchison	10	14	9	13	12	18
Barber	10	41	-	-	7	32
Barton	15	12	12	12	37	36
Bourbon	6	8	8	10	15	25
Brown	-	-	10	22	8	21
Butler	31	11	30	10	47	15
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	15	14	16	16	22	22
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	7	17	11	26	9	24
Cloud	-	-	-	-	7	16
Coffey	6	14	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	29	18	23	15	22	16
Crawford	33	18	38	21	19	11
Decatur	-	-	-	-	-	-
Dickinson	11	13	9	10	12	13
Doniphan	-	-	8	20	-	-
Douglas	43	11	58	14	66	14
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	17	13	24	18	24	19
Ellsworth	-	-	6	22	-	-
Finney	9	9	17	18	15	17
Ford	9	9	8	9	16	21
Franklin	16	14	24	19	18	14
Geary	20	22	19	21	27	23
Gove	-	-	-	-	6	41

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7b. Age-Adjusted Suicide Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	8	19	-	-	7	28
Hamilton	-	-	-	-	-	-
Harper	-	-	7	23	-	-
Harvey	14	9	29	20	20	14
Haskell	6	38	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	10	19
Jefferson	10	11	16	16	12	13
Jewell	-	-	-	-	-	-
Johnson	230	11	247	11	350	15
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	12	12	17	19	9	8
Lane	-	-	-	-	-	-
Leavenworth	43	15	47	15	52	17
Lincoln	6	28	-	-	-	-
Linn	-	-	10	22	11	22
Logan	-	-	-	-	-	-
Lyon	22	16	16	13	27	22
McPherson	15	11	15	11	19	13
Marion	7	9	12	16	7	11
Marshall	-	-	10	20	-	-
Meade	-	-	-	-	-	-
Miami	24	17	25	17	25	16
Mitchell	7	20	7	22	-	-
Montgomery	28	18	21	15	23	16
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	8	14	8	18	9	20
Neosho	17	19	8	11	10	11
Ness	-	-	-	-	-	-
Norton	-	-	-	-	8	30
Osage	9	11	15	20	12	14

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7b. Age-Adjusted Suicide Mortality Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	8	23	-	-	8	26
Phillips	-	-	-	-	6	22
Pottawatomie	11	13	13	13	20	19
Pratt	11	26	9	21	10	26
Rawlins	-	-	-	-	-	-
Reno	41	14	44	16	39	14
Republic	-	-	6	19	-	-
Rice	-	-	10	20	8	15
Riley	22	8	36	16	32	11
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	7	23	9	24	8	23
Saline	27	11	30	13	34	15
Scott	-	-	-	-	-	-
Sedgwick	220	12	265	15	298	16
Seward	8	15	-	-	8	18
Shawnee	124	18	115	16	131	19
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	19	17	19	17	18	16
Thomas	6	19	8	21	-	-
Trego	-	-	-	-	-	-
Wabaunsee	7	19	10	32	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	6	12	-	-	6	15
Woodson	-	-	-	-	-	-
Wyandotte	85	20	77	21	90	25

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7c. Age-Adjusted Suicide Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	45	6	61	7	73	8
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	-	-	-	-	-	-
Semi-Urban	-	-	6	5	8	8
Urban	40	6	51	7	58	8
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7c. Age-Adjusted Suicide Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	-	-	12	9
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7c. Age-Adjusted Suicide Mortality Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	12	7	17	7	17	8
Seward	-	-	-	-	-	-
Shawnee	-	-	-	-	8	9
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	19	9	21	11	14	7

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7d. Age-Adjusted Suicide Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	11	8	12	9	11	8
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	-	-	-	-	-	-
Semi-Urban	-	-	-	-	-	-
Urban	7	9	-	-	-	-
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7d. Age-Adjusted Suicide Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	-	-	-	-
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7d. Age-Adjusted Suicide Mortality Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	-	-	-	-	-	-
Seward	-	-	-	-	-	-
Shawnee	-	-	-	-	-	-
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7e. Age-Adjusted Suicide Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	13	4	14	4	16	4
Frontier	-	-	-	-	-	-
Rural	-	-	-	-	-	-
Dense Rural	-	-	-	-	-	-
Semi-Urban	-	-	-	-	-	-
Urban	7	3	12	4	13	4
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7e. Age-Adjusted Suicide Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	-	-	7	5
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7e. Age-Adjusted Suicide Mortality Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	-	-	-	-
Scott	-	-	-	-	-	-
Sedgwick	-	-	-	-	-	-
Seward	-	-	-	-	-	-
Shawnee	-	-	-	-	-	-
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7f. Age-Adjusted Suicide Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Kansas	45	5	89	7	97	7
Frontier	-	-	-	-	-	-
Rural	-	-	6	12	9	15
Dense Rural	10	4	15	4	24	7
Semi-Urban	-	-	13	10	16	12
Urban	28	6	54	7	43	6
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	6	51
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	-	-	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	6	6	7	9
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7f. Age-Adjusted Suicide Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	-	-	9	5	9	5
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G7f. Age-Adjusted Suicide Mortality Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	Count	Rate	Count	Rate	Count	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	6	25	-	-
Scott	-	-	-	-	-	-
Sedgwick	7	4	23	9	16	6
Seward	-	-	-	-	6	8
Shawnee	7	11	8	10	-	-
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	9	8	11	6	12	6

Note: “-” indicates that data are unavailable or suppressed. Rates are for three five-year periods and age-adjusted per 100,000 persons using the U.S. 2000 standard population. Counts and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8a. All-Cause Mortality Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	876,682	6,543	914,783	6,474	931,829	6,257
Frontier	43,896	7,178	42,562	7,192	42,323	7,503
Rural	79,240	6,477	83,906	6,909	85,486	7,028
Dense Rural	161,907	6,854	164,631	6,829	172,761	7,020
Semi-Urban	135,099	6,444	146,270	6,613	150,014	6,440
Urban	456,314	6,464	476,665	6,227	481,050	5,833
Allen	5,151	7,100	6,363	9,389	6,042	8,827
Anderson	2,998	7,485	2,981	7,487	2,911	7,347
Atchison	5,803	7,096	6,083	7,302	6,056	6,986
Barber	1,827	7,149	1,760	6,763	1,953	8,448
Barton	9,361	6,756	9,260	6,647	10,715	7,712
Bourbon	5,813	7,826	6,945	8,987	6,191	7,936
Brown	3,343	6,381	4,622	9,230	4,724	9,671
Butler	18,219	6,204	20,248	6,295	20,872	6,117
Chase	1,163	7,561	1,276	8,609	872	5,636
Chautauqua	1,466	6,854	1,947	10,056	1,522	8,049
Cherokee	9,023	8,026	11,011	9,785	10,969	9,528
Cheyenne	1,161	8,712	1,166	9,395	913	6,284
Clark	634	5,488	766	7,243	610	5,273
Clay	2,476	6,036	2,751	6,528	2,982	6,901
Cloud	3,245	6,508	3,842	8,176	3,603	7,412
Coffey	2,630	6,000	2,878	6,288	3,608	8,306
Comanche	577	5,265	943	9,845	786	7,589
Cowley	15,331	8,505	14,815	8,007	15,622	8,424
Crawford	13,633	7,404	15,154	7,912	13,657	7,075
Decatur	963	5,534	1,015	6,778	1,426	10,301
Dickinson	6,568	6,963	6,314	6,211	6,841	6,594
Doniphan	2,201	5,350	2,055	4,921	2,254	4,992
Douglas	21,846	5,141	23,293	4,952	22,736	4,465
Edwards	1,714	11,276	776	4,590	1,486	10,681
Elk	1,583	9,824	1,421	9,414	1,277	8,754
Ellis	6,607	4,984	6,927	5,033	7,739	5,335
Ellsworth	2,030	6,430	1,805	5,227	2,009	5,503
Finney	12,014	6,324	10,559	5,860	11,399	5,998
Ford	11,262	6,988	9,720	5,938	11,031	6,200
Franklin	7,800	6,299	9,075	6,889	8,873	6,536
Geary	12,814	9,075	11,705	8,046	13,564	7,691
Gove	762	5,159	830	5,980	1,063	7,653

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8a. All-Cause Mortality Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	967	7,668	1,029	8,145	916	7,174
Grant	2,138	5,475	2,649	6,701	1,752	4,472
Gray	1,665	5,950	2,034	6,855	1,458	4,767
Greeley	346	4,934	365	4,893	514	8,374
Greenwood	3,642	9,932	3,053	8,142	3,039	9,345
Hamilton	1,098	8,339	1,025	7,833	986	7,503
Harper	2,210	6,747	2,841	9,045	2,472	7,915
Harvey	9,313	5,777	10,121	6,035	10,766	6,006
Haskell	1,401	6,730	1,594	6,988	1,463	6,771
Hodgeman	582	5,702	493	4,864	388	3,469
Jackson	4,472	7,042	4,879	7,166	4,176	5,744
Jefferson	6,160	6,517	6,667	6,689	6,766	7,014
Jewell	1,106	5,703	989	5,977	855	5,198
Johnson	96,782	4,211	113,698	4,363	114,682	3,971
Kearny	1,873	8,698	1,205	6,018	1,783	9,329
Kingman	2,449	5,711	2,059	4,659	2,643	6,331
Kiowa	931	5,811	1,090	6,975	941	7,396
Labette	8,914	7,990	9,732	8,681	10,706	9,622
Lane	887	9,663	870	9,635	504	5,737
Leavenworth	21,445	6,208	21,512	5,683	23,983	5,782
Lincoln	1,816	11,177	1,314	7,298	1,024	6,556
Linn	4,115	8,568	3,781	7,378	3,854	7,573
Logan	959	6,383	868	6,158	970	7,177
Lyon	10,612	6,183	10,419	5,935	11,499	6,860
McPherson	7,834	5,418	8,269	5,571	9,221	6,099
Marion	4,495	6,892	4,108	6,589	3,881	6,119
Marshall	2,855	5,231	2,855	5,387	2,960	5,968
Meade	1,536	6,930	1,276	5,760	1,238	5,630
Miami	9,375	6,446	9,450	5,748	10,424	6,061
Mitchell	1,521	4,300	2,082	6,190	2,291	6,961
Montgomery	16,032	9,080	16,873	9,487	14,660	8,053
Morris	1,940	6,437	2,087	7,093	2,070	6,707
Morton	934	5,252	1,156	7,182	1,143	7,122
Nemaha	2,229	4,430	3,169	6,680	3,270	6,496
Neosho	6,281	7,453	6,508	7,749	6,933	8,308
Ness	1,061	6,458	953	5,913	1,173	8,552
Norton	1,558	5,428	1,926	6,339	1,918	6,887
Osage	5,361	6,373	6,156	7,176	6,621	7,726

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8a. All-Cause Mortality Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	1,294	6,100	1,349	6,924	1,458	7,905
Ottawa	2,031	6,331	1,902	5,671	1,963	6,221
Pawnee	2,848	7,806	2,441	6,712	2,903	7,953
Phillips	2,119	7,279	1,631	5,624	2,125	8,036
Pottawatomie	6,036	6,694	5,141	5,006	5,298	4,663
Pratt	3,615	7,794	2,945	6,101	4,091	8,544
Rawlins	650	4,215	1,150	9,896	969	8,182
Reno	20,127	6,149	22,143	6,743	24,374	7,327
Republic	1,877	6,364	1,540	6,685	1,918	8,266
Rice	2,918	5,434	3,552	7,030	3,535	6,889
Riley	10,908	4,498	11,716	4,279	13,395	4,477
Rooks	1,983	7,531	2,080	7,518	1,811	6,927
Rush	1,021	5,798	1,216	7,086	1,303	8,461
Russell	2,499	6,765	2,850	8,004	2,786	7,473
Saline	16,878	6,239	19,785	6,904	19,429	6,667
Scott	1,305	4,956	1,651	6,781	2,174	8,797
Sedgwick	165,966	7,303	177,340	7,314	181,457	6,950
Seward	7,663	6,969	7,776	6,811	6,874	6,060
Shawnee	68,889	7,934	64,731	7,144	65,844	7,001
Sheridan	840	6,059	797	5,867	979	8,926
Sherman	2,438	7,516	2,681	8,857	2,469	8,165
Smith	1,283	5,920	1,277	6,154	1,462	7,737
Stafford	1,257	5,527	1,651	7,464	1,541	6,622
Stanton	1,098	9,182	791	6,985	530	5,046
Stevens	1,710	6,467	2,087	7,436	1,481	5,209
Sumner	9,440	7,470	9,032	7,255	9,808	7,878
Thomas	2,424	6,178	2,885	7,391	2,234	5,505
Trego	1,090	6,655	819	5,550	1,085	6,250
Wabaunsee	2,128	5,948	2,454	6,684	2,263	5,723
Wallace	684	8,029	456	6,471	459	6,091
Washington	2,058	6,834	1,939	6,959	1,654	5,538
Wichita	670	5,247	709	6,432	875	8,331
Wilson	4,694	9,447	4,319	8,474	4,270	8,998
Woodson	1,716	9,712	1,646	8,902	2,220	13,730
Wyandotte	81,386	10,424	76,091	9,597	72,348	8,878

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8b. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	718,520	6,344	722,676	6,210	729,552	6,063
Frontier	40,354	7,003	39,642	7,319	38,104	7,481
Rural	76,008	6,608	76,781	6,800	77,830	7,006
Dense Rural	139,803	6,962	137,907	6,836	143,190	7,150
Semi-Urban	119,982	6,424	126,993	6,568	126,894	6,309
Urban	342,150	6,028	340,992	5,713	343,473	5,421
Allen	4,962	7,160	5,734	8,944	5,715	8,925
Anderson	2,905	7,453	2,898	7,507	2,806	7,398
Atchison	5,215	6,936	5,209	6,719	5,381	6,787
Barber	1,755	7,070	1,663	6,463	1,867	8,472
Barton	8,098	6,493	8,202	6,828	9,656	8,201
Bourbon	5,433	7,722	6,249	8,542	5,932	8,183
Brown	2,778	6,047	3,553	8,089	3,417	8,001
Butler	17,280	6,201	18,655	6,202	18,681	5,891
Chase	1,106	7,423	1,242	8,924	735	4,685
Chautauqua	1,419	7,097	1,879	10,507	1,374	7,877
Cherokee	8,564	8,116	10,015	9,476	10,426	9,821
Cheyenne	1,101	8,428	1,130	9,596	907	6,842
Clark	571	5,038	710	7,168	583	5,644
Clay	2,417	5,997	2,626	6,374	2,822	6,784
Cloud	3,245	6,636	3,808	8,368	3,468	7,442
Coffey	2,528	5,936	2,762	6,313	3,531	8,587
Comanche	563	5,265	890	9,449	673	6,169
Cowley	13,551	8,446	12,836	8,102	12,691	8,129
Crawford	12,809	7,381	14,292	8,058	12,437	7,009
Decatur	957	5,600	978	6,603	1,351	9,952
Dickinson	6,425	7,141	5,909	6,137	6,225	6,372
Doniphan	2,126	5,421	1,946	4,926	2,146	5,127
Douglas	18,856	5,134	18,988	4,704	19,649	4,562
Edwards	1,532	12,019	766	5,280	1,179	11,214
Elk	1,268	7,296	1,414	9,833	1,276	9,200
Ellis	6,415	5,078	6,345	4,888	7,233	5,377
Ellsworth	2,026	7,012	1,646	5,268	1,910	5,907
Finney	6,433	6,243	5,944	6,203	6,333	6,891
Ford	7,835	8,709	5,732	6,898	5,251	6,334
Franklin	7,415	6,319	8,232	6,619	8,228	6,470
Geary	7,845	8,709	7,587	8,139	8,770	7,959
Gove	762	5,211	823	6,087	1,012	7,336

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8b. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	729	5,424	858	7,064	765	6,350
Grant	1,701	6,493	1,496	6,752	1,069	4,726
Gray	1,596	6,285	1,578	6,044	1,335	5,172
Greeley	247	3,327	335	5,499	428	8,399
Greenwood	3,551	9,973	2,989	8,303	2,876	9,366
Hamilton	764	7,352	746	7,373	808	9,103
Harper	2,202	6,916	2,700	9,072	2,148	7,391
Harvey	8,496	5,871	8,985	6,140	9,034	5,820
Haskell	1,164	7,264	1,282	8,217	833	5,413
Hodgeman	577	5,980	493	5,134	388	3,780
Jackson	3,585	6,185	3,825	6,208	3,278	4,972
Jefferson	6,090	6,650	6,302	6,557	6,515	7,051
Jewell	1,106	5,794	973	6,099	776	4,689
Johnson	87,764	4,260	95,468	4,259	96,206	3,939
Kearny	1,044	6,762	930	6,565	1,303	10,029
Kingman	2,346	5,593	2,039	4,771	2,552	6,364
Kiowa	855	5,363	1,078	7,193	866	7,268
Labette	8,060	8,054	8,583	8,596	9,071	9,221
Lane	812	8,808	870	10,422	419	4,794
Leavenworth	17,865	6,137	18,159	5,693	19,891	5,802
Lincoln	1,816	11,358	1,273	7,295	1,024	6,800
Linn	4,043	8,621	3,621	7,299	3,780	7,741
Logan	959	6,507	825	6,158	915	7,069
Lyon	8,699	6,489	8,127	5,872	8,908	7,055
McPherson	7,617	5,483	7,880	5,580	8,548	5,982
Marion	4,415	7,026	3,835	6,411	3,626	5,965
Marshall	2,822	5,245	2,804	5,389	2,746	5,569
Meade	1,503	7,734	1,063	5,441	800	3,974
Miami	8,894	6,379	8,849	5,634	9,929	6,090
Mitchell	1,521	4,349	2,041	6,119	2,134	6,638
Montgomery	13,919	9,118	13,884	9,222	12,731	8,273
Morris	1,920	6,624	2,022	7,308	1,993	6,816
Morton	893	5,981	821	6,087	876	6,926
Nemaha	2,184	4,413	3,106	6,704	3,158	6,465
Neosho	5,845	7,299	6,129	7,727	6,248	8,034
Ness	1,061	6,620	928	6,288	1,054	8,377
Norton	1,499	5,623	1,867	6,693	1,745	6,762
Osage	5,259	6,424	5,947	7,125	6,325	7,720

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8b. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	1,294	6,151	1,346	7,056	1,435	8,003
Ottawa	2,031	6,506	1,827	5,635	1,843	5,952
Pawnee	2,465	7,553	2,216	6,869	2,665	8,653
Phillips	2,119	7,416	1,611	5,771	2,077	8,147
Pottawatomie	5,635	6,522	4,905	5,053	4,905	4,605
Pratt	3,492	8,026	2,635	5,775	3,468	7,802
Rawlins	633	4,184	1,133	10,248	955	8,724
Reno	18,609	6,276	19,510	6,662	20,852	7,117
Republic	1,877	6,470	1,481	6,613	1,902	8,491
Rice	2,666	5,380	3,162	6,865	3,240	7,154
Riley	9,431	4,614	10,119	4,517	10,355	4,226
Rooks	1,921	7,525	2,032	7,650	1,770	6,988
Rush	930	5,265	1,177	7,134	1,274	8,713
Russell	2,470	6,839	2,751	7,970	2,720	7,706
Saline	15,284	6,424	16,880	6,840	15,877	6,441
Scott	1,165	4,724	1,301	6,444	2,043	10,685
Sedgwick	123,763	6,979	123,369	6,752	123,183	6,402
Seward	4,841	8,799	3,637	7,418	3,259	7,169
Shawnee	53,364	7,535	48,261	6,696	49,465	6,665
Sheridan	840	6,166	797	6,095	784	7,355
Sherman	2,098	7,192	2,389	9,088	2,214	8,501
Smith	1,283	5,987	1,247	6,168	1,462	7,972
Stafford	1,257	6,025	1,553	7,972	1,424	6,940
Stanton	975	11,303	602	8,782	363	5,927
Stevens	1,495	7,392	1,591	8,114	938	4,712
Sumner	9,115	7,684	8,451	7,280	9,144	7,912
Thomas	2,424	6,375	2,695	7,179	1,980	5,140
Trego	1,090	6,740	809	5,685	1,066	6,352
Wabaunsee	2,102	6,056	2,364	6,664	2,158	5,669
Wallace	610	7,322	434	7,072	426	5,795
Washington	2,058	6,936	1,869	6,930	1,582	5,490
Wichita	472	4,571	446	5,298	676	8,654
Wilson	4,616	9,591	4,012	8,094	3,987	8,825
Woodson	1,716	10,009	1,582	8,770	2,154	14,035
Wyandotte	40,538	9,644	36,747	9,322	35,079	9,081

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8c. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	93,233	11,993	93,989	11,022	89,573	9,663
Frontier	343	16,624	231	7,200	240	6,989
Rural	510	5,753	582	5,858	769	5,604
Dense Rural	3,947	10,214	4,867	11,483	4,806	9,626
Semi-Urban	8,904	9,882	8,793	9,870	8,980	8,842
Urban	79,529	12,542	79,456	11,358	74,694	9,949
Allen	119	12,176	291	23,045	78	7,098
Anderson	65	16,896	-	-	20	5,816
Atchison	471	11,831	652	14,092	367	6,850
Barber	15	11,897	-	-	-	-
Barton	231	12,996	153	8,428	46	2,883
Bourbon	332	14,462	425	22,125	70	3,116
Brown	28	3,140	48	4,840	92	6,752
Butler	364	7,597	413	10,269	373	7,416
Chase	57	35,911	-	-	-	-
Chautauqua	4	6,604	-	-	4	2,641
Cherokee	102	8,681	314	46,992	102	14,419
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	59	15,731	8	10,566	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	24	10,787	-	-
Comanche	-	-	8	6,604	-	-
Cowley	394	8,565	245	5,067	782	14,369
Crawford	352	11,387	327	9,008	385	11,113
Decatur	-	-	-	-	-	-
Dickinson	40	7,071	176	21,003	132	10,530
Doniphan	54	7,711	68	9,725	22	2,431
Douglas	1,775	9,563	1,554	7,307	1,463	6,102
Edwards	34	34,694	-	-	25	37,454
Elk	-	-	-	-	-	-
Ellis	-	-	161	27,018	106	6,290
Ellsworth	-	-	-	-	-	-
Finney	44	4,308	142	4,930	264	5,009
Ford	185	10,836	140	5,207	184	4,065
Franklin	45	3,537	197	10,820	46	2,272
Geary	3,650	11,565	2,698	9,666	2,973	8,956
Gove	-	-	-	-	51	176,774

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8c. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	178	58,953	163	40,523	85	20,255
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	18	8,725	-	-
Greenwood	-	-	-	-	-	-
Hamilton	8	6,604	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	175	6,897	365	9,575	280	9,033
Haskell	70	263,739	-	-	-	-
Hodgeman	5	3,302	-	-	-	-
Jackson	157	42,747	38	9,507	141	22,100
Jefferson	62	16,116	68	15,455	20	4,892
Jewell	-	-	-	-	-	-
Johnson	4,298	6,931	7,221	6,939	8,097	5,949
Kearny	-	-	12	8,725	-	-
Kingman	16	27,919	-	-	41	24,167
Kiowa	-	-	-	-	-	-
Labette	737	14,700	653	12,416	1,092	17,075
Lane	-	-	-	-	-	-
Leavenworth	3,000	8,948	2,530	7,335	2,561	6,755
Lincoln	-	-	-	-	-	-
Linn	-	-	28	10,487	15	2,974
Logan	-	-	-	-	-	-
Lyon	410	9,946	344	7,307	256	8,585
McPherson	47	6,386	82	9,115	213	10,863
Marion	2	550	47	16,169	-	-
Marshall	-	-	3	1,415	75	17,273
Meade	-	-	-	-	-	-
Miami	329	15,475	385	16,130	209	7,365
Mitchell	-	-	-	-	36	24,392
Montgomery	1,522	13,981	1,847	19,695	1,115	11,299
Morris	-	-	-	-	52	12,016
Morton	-	-	-	-	-	-
Nemaha	-	-	3	2,201	-	-
Neosho	125	25,445	123	12,874	119	8,475
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	11	6,398	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8c. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	23	23,855
Ottawa	-	-	-	-	-	-
Pawnee	188	11,697	170	9,517	84	4,712
Phillips	-	-	20	34,899	-	-
Pottawatomie	47	7,544	47	5,180	30	3,410
Pratt	54	15,382	32	5,903	48	10,231
Rawlins	-	-	-	-	7	23,113
Reno	740	9,707	686	7,818	962	10,221
Republic	-	-	-	-	-	-
Rice	4	1,101	19	7,535	34	10,052
Riley	789	5,062	739	4,479	1,594	7,428
Rooks	10	4,717	-	-	-	-
Rush	-	-	-	-	-	-
Russell	3	825	90	17,439	10	3,885
Saline	938	9,872	1,136	10,979	1,043	8,701
Scott	-	-	-	-	-	-
Sedgwick	27,375	12,938	29,417	12,448	29,081	11,480
Seward	353	9,957	644	17,327	651	17,251
Shawnee	10,385	13,337	9,948	11,985	9,177	11,130
Sheridan	-	-	-	-	30	67,417
Sherman	32	16,786	30	15,558	15	6,232
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	27	14,616
Sumner	26	3,052	101	10,515	131	13,330
Thomas	-	-	75	25,909	100	24,436
Trego	-	-	-	-	-	-
Wabaunsee	13	7,089	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	57	23,948
Wichita	-	-	-	-	-	-
Wilson	8	3,108	15	10,906	63	20,688
Woodson	-	-	-	-	15	32,718
Wyandotte	32,696	14,840	28,786	13,306	24,315	11,418

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: *Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.*

Figure G8d. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	9,548	7,652	10,703	8,252	11,096	7,708
Frontier	316	11,645	252	6,606	309	9,306
Rural	1,029	11,342	1,368	14,507	1,727	15,998
Dense Rural	2,147	8,303	2,416	8,930	2,509	7,966
Semi-Urban	1,223	5,757	1,470	6,885	1,292	5,459
Urban	4,833	7,354	5,195	7,650	5,259	7,196
Allen	49	8,363	45	11,239	18	3,569
Anderson	28	18,402	52	23,245	-	-
Atchison	94	27,328	99	20,540	31	4,508
Barber	-	-	-	-	-	-
Barton	72	13,072	69	11,229	87	13,042
Bourbon	-	-	30	4,445	-	-
Brown	537	12,769	725	18,423	969	22,133
Butler	188	6,006	233	7,273	106	2,857
Chase	-	-	11	95,972	-	-
Chautauqua	43	5,509	39	4,064	141	21,255
Cherokee	247	6,013	190	4,196	241	4,320
Cheyenne	-	-	-	-	-	-
Clark	63	51,499	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	23	37,971	56	27,073	-	-
Comanche	14	8,725	-	-	30	67,417
Cowley	485	12,627	419	10,611	525	12,488
Crawford	128	6,858	93	4,509	96	3,082
Decatur	-	-	37	46,282	-	-
Dickinson	45	12,994	-	-	59	11,055
Doniphan	18	2,855	14	2,255	36	6,037
Douglas	839	6,303	738	7,248	465	4,494
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	25	9,631	59	25,311	-	-
Ellsworth	-	-	-	-	37	12,214
Finney	-	-	29	2,715	43	4,180
Ford	-	-	74	7,048	100	8,908
Franklin	16	1,702	211	21,830	177	14,112
Geary	252	10,604	18	1,540	70	3,417
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8d. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	47	22,633
Gray	-	-	47	35,400	28	34,321
Greeley	-	-	-	-	-	-
Greenwood	29	7,890	-	-	15	1,925
Hamilton	-	-	-	-	-	-
Harper	6	6,604	66	28,813	86	30,042
Harvey	19	3,226	99	11,581	90	8,078
Haskell	-	-	-	-	24	26,967
Hodgeman	-	-	-	-	-	-
Jackson	626	14,986	699	14,662	605	11,187
Jefferson	8	5,283	117	10,428	77	6,677
Jewell	-	-	-	-	46	22,078
Johnson	573	5,793	823	8,071	372	3,463
Kearny	-	-	-	-	18	6,828
Kingman	74	24,207	-	-	17	4,635
Kiowa	46	623,636	12	14,957	-	-
Labette	96	3,681	100	4,171	133	5,745
Lane	-	-	-	-	16	9,971
Leavenworth	96	3,234	92	3,677	157	4,525
Lincoln	-	-	41	53,916	-	-
Linn	72	22,237	18	3,414	-	-
Logan	-	-	19	15,070	-	-
Lyon	94	12,503	61	11,985	170	13,702
McPherson	-	-	60	9,907	39	4,716
Marion	41	14,931	31	11,322	46	7,441
Marshall	-	-	13	10,311	-	-
Meade	22	22,020	15	5,453	6	2,476
Miami	-	-	37	3,141	29	2,556
Mitchell	-	-	41	79,407	12	4,986
Montgomery	116	1,874	297	3,977	227	3,589
Morris	13	8,725	16	46,532	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	112	32,439
Neosho	87	8,962	171	16,162	86	9,927
Ness	-	-	-	-	-	-
Norton	54	31,195	30	19,262	-	-
Osage	17	5,613	22	4,686	117	12,309

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8d. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	7	3,302	23	6,329
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	21	4,230	37	4,169
Pratt	30	40,450	104	59,211	72	76,030
Rawlins	17	37,080	-	-	-	-
Reno	215	10,313	201	8,976	234	8,622
Republic	-	-	-	-	16	11,366
Rice	67	25,831	-	-	42	11,165
Riley	127	8,846	70	9,672	60	5,853
Rooks	52	65,542	23	25,843	-	-
Rush	19	65,200	-	-	-	-
Russell	-	-	-	-	-	-
Saline	162	11,521	211	17,277	203	13,841
Scott	-	-	-	-	-	-
Sedgwick	2,010	7,946	2,076	7,709	2,375	8,285
Seward	88	13,081	74	13,896	52	10,713
Shawnee	954	9,425	1,044	9,922	1,434	12,703
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	33	28,747
Smith	-	-	-	-	-	-
Stafford	-	-	6	2,641	-	-
Stanton	11	18,160	-	-	-	-
Stevens	12	20,939	50	21,068	18	5,415
Sumner	95	6,848	63	5,504	53	2,298
Thomas	-	-	9	19,811	-	-
Trego	-	-	10	6,604	-	-
Wabaunsee	-	-	41	20,876	29	9,775
Wallace	-	-	22	42,376	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	65	14,417	48	8,192	96	13,354
Woodson	-	-	-	-	46	172,546
Wyandotte	361	5,958	422	6,390	456	6,917

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8e. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	7,257	2,799	8,907	2,839	10,295	2,669
Frontier	30	2,452	23	981	60	2,287
Rural	110	3,172	132	3,172	301	6,218
Dense Rural	1,182	4,753	600	2,364	1,005	3,399
Semi-Urban	1,214	4,328	896	2,525	1,347	3,219
Urban	4,721	2,352	7,256	2,915	7,540	2,471
Allen	8	6,604	-	-	14	3,054
Anderson	-	-	18	11,217	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	39	12,684	58	17,253	-	-
Bourbon	-	-	30	13,744	-	-
Brown	-	-	-	-	41	27,119
Butler	-	-	78	3,308	57	2,010
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	90	27,351	7	3,082	66	5,906
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	10	3,885
Cloud	-	-	-	-	67	28,685
Coffey	18	8,725	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	122	5,593	10	635	222	7,163
Crawford	219	9,269	37	1,829	93	2,984
Decatur	-	-	-	-	-	-
Dickinson	26	7,664	12	3,173	-	-
Doniphan	-	-	-	-	-	-
Douglas	55	99	271	2,363	258	1,365
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	48	2,109	64	2,232	11	835
Ellsworth	-	-	-	-	-	-
Finney	360	5,763	89	1,647	159	2,695
Ford	71	2,501	36	1,452	49	1,433
Franklin	50	14,122	-	-	68	9,432
Geary	540	8,561	261	3,336	452	4,465
Gove	-	-	7	6,604	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansas. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8e. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	70	6,792	61	4,575	143	7,752
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	81	39,713	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	981	1,328	2,412	2,194	2,042	1,605
Kearny	-	-	-	-	-	-
Kingman	13	26,634	-	-	-	-
Kiowa	30	101,126	-	-	-	-
Labette	-	-	-	-	48	9,305
Lane	-	-	-	-	-	-
Leavenworth	202	3,821	46	837	162	2,262
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	114	3,421	145	5,675	308	6,610
McPherson	42	8,656	24	3,517	-	-
Marion	-	-	-	-	24	15,268
Marshall	-	-	-	-	-	-
Meade	-	-	16	34,899	31	33,607
Miami	-	-	-	-	31	4,132
Mitchell	-	-	-	-	-	-
Montgomery	34	4,463	57	5,558	-	-
Morris	7	5,778	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	60	277,292	-	-
Neosho	36	15,822	49	9,100	71	18,429
Ness	-	-	-	-	-	-
Norton	-	-	-	-	71	78,550
Osage	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8e. All-Cause Mortality Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	70	51,545	-	-	-	-
Phillips	-	-	-	-	7	2,568
Pottawatomie	109	34,104	-	-	-	-
Pratt	-	-	10	5,503	-	-
Rawlins	-	-	-	-	-	-
Reno	87	5,784	63	3,991	32	1,758
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	120	1,764	195	1,727	219	1,832
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	90	1,794	139	2,217	251	3,755
Scott	-	-	40	25,017	-	-
Sedgwick	2,665	3,175	3,872	4,031	3,670	3,182
Seward	34	1,460	66	1,771	43	1,020
Shawnee	165	1,694	244	2,114	185	1,351
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	10	3,302
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	14	2,841
Thomas	-	-	-	-	28	10,204
Trego	-	-	-	-	19	11,841
Wabaunsee	-	-	-	-	10	6,003
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	43	16,892
Woodson	-	-	-	-	-	-
Wyandotte	653	4,955	411	3,403	1,223	5,821

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8f. All-Cause Mortality Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	46,909	4,818	61,777	4,885	67,549	4,589
Frontier	2,847	7,524	2,030	5,732	2,809	6,334
Rural	1,574	3,738	3,962	6,369	3,557	4,990
Dense Rural	14,708	5,368	15,993	4,963	17,929	4,955
Semi-Urban	3,446	3,861	5,613	4,457	7,474	5,203
Urban	24,334	4,684	34,056	4,720	35,772	4,230
Allen	13	1,274	265	10,853	29	1,872
Anderson	-	-	-	-	15	3,966
Atchison	23	1,519	48	4,955	41	3,238
Barber	57	10,389	83	16,975	-	-
Barton	921	6,301	708	5,448	626	3,729
Bourbon	9	2,377	68	8,033	77	12,516
Brown	-	-	175	9,406	108	10,905
Butler	383	5,127	345	3,881	713	6,376
Chase	-	-	-	-	58	7,116
Chautauqua	-	-	6	3,048	-	-
Cherokee	20	2,743	210	16,963	14	1,001
Cheyenne	60	21,264	36	9,757	6	1,467
Clark	-	-	56	6,114	21	6,292
Clay	-	-	117	12,115	-	-
Cloud	-	-	34	4,808	68	6,790
Coffey	61	10,681	19	4,876	41	5,939
Comanche	-	-	-	-	35	10,540
Cowley	779	9,512	955	7,135	851	5,741
Crawford	125	1,299	275	3,969	371	4,936
Decatur	-	-	-	-	75	51,818
Dickinson	32	1,808	113	4,660	272	5,450
Doniphan	-	-	-	-	50	13,973
Douglas	296	2,120	871	4,519	542	2,540
Edwards	148	6,038	10	1,001	258	9,050
Elk	315	67,325	-	-	-	-
Ellis	63	1,742	216	3,104	300	5,500
Ellsworth	-	-	97	6,711	62	3,288
Finney	5,176	6,440	4,177	5,718	4,403	5,139
Ford	3,167	5,009	3,550	4,063	5,193	5,199
Franklin	255	6,786	170	5,701	197	4,745
Geary	454	4,950	807	5,003	737	3,554
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansas. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G8f. All-Cause Mortality Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	60	28,685	8	3,774	42	15,672
Grant	437	3,149	1,104	6,637	623	4,210
Gray	69	5,892	409	9,462	95	1,975
Greeley	99	9,030	12	4,188	86	11,337
Greenwood	62	7,844	-	-	81	9,002
Hamilton	326	12,882	221	6,060	151	6,201
Harper	-	-	75	2,657	191	13,735
Harvey	553	4,056	407	2,445	1,082	5,717
Haskell	167	4,121	213	5,660	584	8,624
Hodgeman	-	-	-	-	-	-
Jackson	-	-	96	4,402	83	2,823
Jefferson	-	-	57	4,803	75	4,530
Jewell	-	-	-	-	-	-
Johnson	3,051	3,260	5,808	3,440	5,494	2,747
Kearny	829	12,808	263	5,238	461	7,791
Kingman	-	-	20	3,242	-	-
Kiowa	-	-	-	-	75	10,364
Labette	21	1,076	140	5,563	151	4,590
Lane	75	11,515	-	-	69	16,261
Leavenworth	282	2,314	473	3,585	821	4,388
Lincoln	-	-	-	-	-	-
Linn	-	-	71	10,008	11	1,714
Logan	-	-	24	9,408	52	7,670
Lyon	1,295	4,561	1,537	4,164	1,669	4,576
McPherson	128	4,210	123	4,451	409	8,590
Marion	37	6,188	155	14,603	99	4,474
Marshall	33	11,925	35	5,526	139	19,342
Meade	11	3,027	107	3,654	401	9,479
Miami	77	1,600	155	4,962	130	4,109
Mitchell	-	-	-	-	90	8,936
Montgomery	366	8,648	524	9,255	322	4,010
Morris	-	-	-	-	-	-
Morton	41	2,293	332	12,364	210	8,118
Nemaha	34	11,518	-	-	-	-
Neosho	188	8,612	7	284	353	12,490
Ness	-	-	25	6,242	44	2,286
Norton	-	-	20	5,816	-	-
Osage	74	8,532	111	7,696	35	2,145

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: *Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.*

Figure G8f. All-Cause Mortality Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	52	21,083	105	18,313
Pawnee	125	8,474	21	1,656	95	5,597
Phillips	-	-	-	-	36	10,841
Pottawatomie	245	8,155	144	4,128	248	5,770
Pratt	39	5,066	144	7,528	386	10,212
Rawlins	-	-	17	7,806	-	-
Reno	401	2,634	1,305	5,070	1,698	6,578
Republic	-	-	-	-	-	-
Rice	181	6,362	272	8,616	164	5,779
Riley	428	5,963	395	2,351	725	3,792
Rooks	-	-	7	2,889	41	14,628
Rush	72	398,282	12	5,816	29	8,413
Russell	23	8,867	9	11,887	49	9,059
Saline	404	1,852	1,230	5,602	1,499	6,070
Scott	140	10,520	282	8,460	131	3,798
Sedgwick	9,914	5,053	13,912	5,388	15,638	4,921
Seward	2,347	4,844	3,296	4,846	2,735	4,277
Shawnee	3,745	5,770	4,023	5,027	4,067	4,340
Sheridan	-	-	-	-	148	30,450
Sherman	308	9,632	235	8,930	108	3,877
Smith	-	-	30	12,641	-	-
Stafford	-	-	92	5,324	64	2,387
Stanton	112	3,154	189	5,027	167	4,736
Stevens	203	3,120	446	5,678	331	4,163
Sumner	204	3,548	172	3,764	315	7,227
Thomas	-	-	31	2,948	108	3,645
Trego	-	-	-	-	-	-
Wabaunsee	9	2,049	49	5,499	26	2,988
Wallace	74	20,951	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	198	6,745	263	11,910	124	4,969
Wilson	-	-	112	13,961	-	-
Woodson	-	-	-	-	-	-
Wyandotte	7,046	5,191	8,969	4,909	9,210	4,789

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9a. Coronary Heart Disease Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	83,219	628	80,542	557	75,176	480
Frontier	4,868	663	3,859	535	3,715	506
Rural	8,671	641	8,412	606	7,109	518
Dense Rural	15,475	657	15,996	644	14,913	565
Semi-Urban	13,484	660	12,515	565	12,198	524
Urban	40,686	601	39,660	518	37,233	435
Allen	617	842	684	999	490	625
Anderson	379	853	273	590	185	503
Atchison	568	694	588	725	384	380
Barber	173	490	116	331	77	225
Barton	967	628	1,269	832	1,235	803
Bourbon	663	817	690	864	740	871
Brown	358	630	587	978	435	698
Butler	1,753	587	1,797	525	1,721	468
Chase	167	990	83	451	104	566
Chautauqua	216	802	162	686	114	396
Cherokee	1,462	1,272	1,323	1,091	1,040	797
Cheyenne	120	642	127	639	26	110
Clark	80	528	60	360	115	770
Clay	140	311	180	388	127	388
Cloud	365	662	309	656	377	700
Coffey	295	612	287	559	483	880
Comanche	35	232	101	628	92	890
Cowley	1,319	697	1,386	675	1,577	773
Crawford	1,873	1,047	1,572	849	1,358	699
Decatur	79	350	205	985	75	339
Dickinson	700	655	763	681	752	628
Doniphan	320	755	246	521	341	731
Douglas	2,182	569	1,777	388	1,598	329
Edwards	71	346	62	291	130	629
Elk	211	1,070	203	1,229	203	977
Ellis	350	264	444	352	643	411
Ellsworth	277	761	250	588	221	630
Finney	792	541	733	446	823	468
Ford	909	703	756	531	620	383
Franklin	570	455	982	722	702	506
Geary	1,054	950	783	643	1,005	739
Gove	75	435	72	378	53	307

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9a. Coronary Heart Disease Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	102	503	76	324	89	491
Grant	246	697	248	691	115	262
Gray	151	571	233	765	66	204
Greeley	33	433	31	369	13	128
Greenwood	487	1,061	291	671	331	799
Hamilton	70	552	84	699	89	764
Harper	345	919	320	908	114	302
Harvey	953	558	592	324	759	388
Haskell	186	973	151	727	106	516
Hodgeman	24	191	9	61	39	265
Jackson	502	732	446	567	406	479
Jefferson	431	411	677	583	452	347
Jewell	193	724	104	377	113	445
Johnson	8,393	376	8,381	312	8,918	288
Kearny	126	641	37	187	156	793
Kingman	161	306	276	564	286	624
Kiowa	81	410	118	661	37	195
Labette	1,105	944	976	829	620	484
Lane	101	1,090	155	1,658	116	1,337
Leavenworth	2,373	691	3,015	766	2,917	642
Lincoln	242	1,145	123	655	118	497
Linn	414	721	563	939	636	1,110
Logan	110	682	20	102	85	407
Lyon	1,052	670	1,006	609	976	570
McPherson	749	503	734	467	838	515
Marion	362	449	192	228	261	402
Marshall	274	437	289	432	275	380
Meade	129	541	132	556	72	271
Miami	920	608	866	490	1,151	596
Mitchell	162	399	203	588	204	487
Montgomery	1,521	777	1,380	722	1,164	534
Morris	280	869	112	283	195	441
Morton	171	972	124	765	150	721
Nemaha	245	438	352	716	231	362
Neosho	516	571	689	713	633	693
Ness	127	659	95	704	90	427
Norton	125	373	178	592	77	194
Osage	542	565	797	788	508	442

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9a. Coronary Heart Disease Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	182	723	104	398	141	518
Ottawa	161	454	189	484	212	591
Pawnee	327	801	314	803	171	369
Phillips	135	422	184	641	46	104
Pottawatomie	510	560	519	505	539	489
Pratt	205	370	266	455	314	677
Rawlins	83	412	140	875	26	186
Reno	1,954	571	1,679	455	1,980	573
Republic	367	1,155	126	338	194	601
Rice	355	620	298	477	381	773
Riley	859	447	773	367	590	279
Rooks	164	488	113	340	116	314
Rush	145	883	175	793	119	582
Russell	334	809	343	686	270	704
Saline	2,027	736	2,091	699	1,768	597
Scott	107	380	75	248	170	647
Sedgwick	14,927	693	14,420	597	13,791	522
Seward	559	645	316	351	392	388
Shawnee	7,585	843	7,275	755	5,701	550
Sheridan	92	550	104	578	107	852
Sherman	390	1,066	270	694	231	692
Smith	152	497	110	379	192	686
Stafford	168	609	169	648	98	292
Stanton	82	725	26	217	43	298
Stevens	269	1,064	165	596	88	301
Sumner	842	616	954	699	904	654
Thomas	189	460	236	538	73	144
Trego	118	599	54	235	67	243
Wabaunsee	250	624	228	561	210	585
Wallace	55	542	-	-	36	320
Washington	237	656	184	508	91	323
Wichita	14	92	4	27	52	387
Wilson	609	1,093	594	1,051	327	491
Woodson	361	1,840	207	839	168	735
Wyandotte	5,226	718	4,792	635	4,308	531

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9b. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	74,448	630	70,047	557	64,009	479
Frontier	4,799	678	3,610	526	3,517	505
Rural	8,382	641	8,023	602	6,761	522
Dense Rural	14,365	669	14,534	656	13,294	582
Semi-Urban	12,485	660	11,376	563	10,960	524
Urban	34,382	597	32,418	513	29,477	423
Allen	574	820	611	922	458	623
Anderson	379	864	273	601	161	459
Atchison	491	636	510	669	365	389
Barber	173	496	116	337	53	137
Barton	929	631	1,214	870	1,141	815
Bourbon	645	831	550	699	705	877
Brown	315	612	470	873	396	696
Butler	1,722	600	1,690	516	1,621	464
Chase	167	1,022	62	325	104	585
Chautauqua	216	832	162	726	103	388
Cherokee	1,413	1,293	1,185	1,017	943	752
Cheyenne	118	642	127	654	26	113
Clark	74	500	60	368	115	795
Clay	140	315	180	396	127	404
Cloud	365	668	275	566	363	699
Coffey	277	583	275	553	454	853
Comanche	35	233	101	637	92	930
Cowley	1,284	738	1,279	684	1,364	759
Crawford	1,858	1,075	1,466	826	1,350	739
Decatur	79	351	205	996	75	343
Dickinson	663	639	692	629	752	654
Doniphan	302	738	199	436	311	709
Douglas	2,051	590	1,500	363	1,389	324
Edwards	71	365	62	308	130	699
Elk	211	1,093	201	1,260	203	1,012
Ellis	350	268	394	327	580	392
Ellsworth	277	805	250	619	221	702
Finney	711	696	560	508	585	600
Ford	764	732	722	744	486	445
Franklin	554	460	952	734	639	475
Geary	682	862	565	682	733	791
Gove	75	438	65	352	53	312

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9b. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	100	520	61	286	81	489
Grant	196	720	158	535	88	262
Gray	151	596	233	813	66	219
Greeley	33	461	1	12	13	136
Greenwood	478	1,068	291	691	331	832
Hamilton	70	614	84	861	86	1,031
Harper	345	934	320	937	114	313
Harvey	886	556	523	316	660	379
Haskell	116	710	144	885	106	686
Hodgeman	24	192	9	62	39	274
Jackson	456	717	344	472	339	444
Jefferson	431	420	635	563	432	344
Jewell	193	728	88	321	99	409
Johnson	7,993	387	7,491	311	7,853	286
Kearny	124	720	26	156	76	447
Kingman	161	309	276	576	245	490
Kiowa	81	413	118	680	37	200
Labette	965	898	876	820	597	520
Lane	101	1,115	155	1,748	116	1,426
Leavenworth	2,091	692	2,598	755	2,511	651
Lincoln	242	1,159	111	623	118	510
Linn	412	729	563	964	621	1,134
Logan	110	695	20	104	85	418
Lyon	965	706	926	673	844	630
McPherson	720	496	693	458	823	528
Marion	360	454	192	232	251	405
Marshall	274	441	289	437	275	386
Meade	129	560	132	596	72	296
Miami	867	593	804	460	1,144	617
Mitchell	162	401	162	389	204	493
Montgomery	1,393	795	1,231	738	1,064	553
Morris	280	891	112	291	195	456
Morton	165	1,053	86	550	138	782
Nemaha	245	443	352	732	231	369
Neosho	500	575	683	736	524	599
Ness	127	666	95	806	90	437
Norton	125	384	178	664	77	198
Osage	530	562	797	805	485	434

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9b. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	182	727	104	402	118	419
Ottawa	161	459	189	492	205	588
Pawnee	280	751	312	895	153	369
Phillips	135	427	184	654	46	105
Pottawatomie	470	529	519	524	417	408
Pratt	196	366	250	446	274	576
Rawlins	66	324	140	895	26	190
Reno	1,867	586	1,611	472	1,790	575
Republic	367	1,165	126	342	190	601
Rice	352	644	298	501	347	733
Riley	765	442	708	387	470	258
Rooks	164	494	106	327	116	320
Rush	145	898	148	671	96	471
Russell	331	818	334	683	270	734
Saline	1,891	741	1,826	674	1,489	566
Scott	107	395	75	260	170	738
Sedgwick	12,719	707	11,798	604	10,484	501
Seward	364	596	299	531	306	553
Shawnee	6,305	820	5,864	734	4,486	521
Sheridan	92	557	104	583	107	901
Sherman	365	1,041	261	713	231	772
Smith	152	500	80	260	192	695
Stafford	168	631	163	675	98	304
Stanton	82	844	8	68	43	389
Stevens	227	1,025	129	558	51	221
Sumner	838	641	846	662	837	651
Thomas	189	466	236	551	73	148
Trego	118	604	54	237	67	246
Wabaunsee	250	637	228	578	205	598
Wallace	55	561	-	-	36	330
Washington	237	664	184	515	91	334
Wichita	14	100	4	31	52	462
Wilson	609	1,121	569	1,036	323	501
Woodson	361	1,883	207	854	168	753
Wyandotte	3,223	717	3,167	766	2,754	626

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9c. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	5,756	940	5,648	780	5,492	675
Frontier	2	72	25	1,057	31	963
Rural	95	1,169	24	368	100	920
Dense Rural	370	1,189	499	1,428	332	827
Semi-Urban	571	982	602	874	455	598
Urban	4,718	919	4,498	740	4,574	672
Allen	7	405	45	5,869	-	-
Anderson	-	-	-	-	-	-
Atchison	73	2,114	76	1,845	19	407
Barber	-	-	-	-	-	-
Barton	29	2,813	-	-	6	357
Bourbon	18	730	97	6,060	35	1,555
Brown	8	574	13	1,334	-	-
Butler	-	-	27	552	16	421
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	8	839	19	4,144	25	3,210
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	30	965	13	275	68	1,307
Crawford	15	1,073	68	2,932	8	452
Decatur	-	-	-	-	-	-
Dickinson	-	-	37	5,957	-	-
Doniphan	-	-	45	6,241	10	1,139
Douglas	119	948	114	805	102	515
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	24	4,558	3	991
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	45	1,361
Ford	18	1,848	2	159	-	-
Franklin	-	-	-	-	-	-
Geary	243	1,252	130	694	192	823
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander. Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9c. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	2	118	7	825	8	1,321
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	18	8,725	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	24	1,051	48	1,590	15	469
Haskell	70	263,739	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	15	2,045	-	-
Jewell	-	-	-	-	-	-
Johnson	292	665	471	587	405	360
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	41	24,167
Kiowa	-	-	-	-	-	-
Labette	127	2,728	47	943	16	242
Lane	-	-	-	-	-	-
Leavenworth	242	887	336	1,090	263	733
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	15	2,974
Logan	-	-	-	-	-	-
Lyon	8	518	6	392	39	1,487
McPherson	9	2,972	19	2,810	9	1,213
Marion	2	550	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	53	1,878	20	772	-	-
Mitchell	-	-	-	-	-	-
Montgomery	59	651	116	1,218	61	565
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	4	600	-	-	21	2,360
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	11	6,398	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander. Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9c. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	23	23,855
Ottawa	-	-	-	-	-	-
Pawnee	-	-	2	183	10	805
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	23	1,632
Pratt	9	1,121	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	71	1,030	2	35	109	1,280
Republic	-	-	-	-	-	-
Rice	3	825	-	-	34	10,052
Riley	19	366	38	463	3	54
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	3	825	9	4,245	-	-
Saline	87	1,230	153	1,818	51	552
Scott	-	-	-	-	-	-
Sedgwick	1,550	948	1,340	676	1,866	847
Seward	28	860	4	129	2	66
Shawnee	967	1,475	937	1,250	765	959
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	50	5,740	11	979
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	1,548	795	1,300	618	1,173	540

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. "-" indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander. Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9d. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	706	694	1,065	849	1,087	778
Frontier	27	1,164	18	470	25	488
Rural	59	850	211	2,241	44	440
Dense Rural	166	756	217	787	193	600
Semi-Urban	145	798	180	909	140	654
Urban	309	581	437	680	685	950
Allen	23	3,926	-	-	18	3,569
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	9	2,702	25	3,391	-	-
Bourbon	-	-	-	-	-	-
Brown	35	1,133	100	2,970	39	1,085
Butler	27	755	7	335	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	11	1,038
Cherokee	4	131	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	6	9,906	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	10	16,509	12	6,980	-	-
Comanche	-	-	-	-	-	-
Cowley	5	201	39	927	33	873
Crawford	-	-	38	2,191	-	-
Decatur	-	-	-	-	-	-
Dickinson	13	2,315	-	-	-	-
Doniphan	18	2,855	2	322	18	3,205
Douglas	7	203	103	1,139	69	770
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	29	2,715	-	-
Ford	-	-	-	-	-	-
Franklin	16	1,702	13	1,990	31	2,334
Geary	-	-	18	1,540	-	-
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander. Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9d. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	4	1,761	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	1	206	-	-	24	2,501
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	46	1,201	77	1,624	67	1,163
Jefferson	-	-	27	2,304	-	-
Jewell	-	-	-	-	14	5,816
Johnson	-	-	33	294	41	455
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	12	446	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	-	-	47	1,404
Lincoln	-	-	12	10,470	-	-
Linn	2	2,641	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	25	4,372	-	-	-	-
McPherson	-	-	18	3,363	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	41	79,407	-	-
Montgomery	4	70	28	401	15	200
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	1	330	-	-	14	1,259

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander. Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9d. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	30	2,953
Pratt	-	-	6	6,604	1	3,302
Rawlins	17	37,080	-	-	-	-
Reno	-	-	54	2,382	-	-
Republic	-	-	-	-	4	2,641
Rice	-	-	-	-	-	-
Riley	48	4,697	-	-	23	1,615
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	49	3,398	22	2,494	47	3,483
Scott	-	-	-	-	-	-
Sedgwick	141	627	76	340	335	1,145
Seward	6	3,048	-	-	-	-
Shawnee	119	1,376	199	1,897	173	1,463
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	6	2,641	-	-
Stanton	-	-	-	-	-	-
Stevens	12	20,939	27	11,377	-	-
Sumner	4	498	-	-	13	459
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	25	4,267	-	-
Woodson	-	-	-	-	-	-
Wyandotte	42	603	26	472	20	269

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander. Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9e. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	442	226	523	194	522	151
Frontier	-	-	7	392	-	-
Rural	-	-	10	317	9	188
Dense Rural	118	513	37	181	57	203
Semi-Urban	114	538	63	176	62	187
Urban	210	141	406	191	394	142
Allen	-	-	-	-	14	3,054
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	30	13,744	-	-
Brown	-	-	-	-	-	-
Butler	4	629	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	37	8,980	-	-	1	206
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	2	127	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	24	7,035	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	23	314	28	240
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	20	314	-	-	-	-
Ford	19	650	-	-	2	78
Franklin	-	-	-	-	-	-
Geary	69	1,013	52	653	16	162
Gove	-	-	7	6,604	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander. Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9e. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	49	101	111	134	121	98
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	16	333	3	65	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	15	509
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	34	4,463	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	25	9,110
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander. Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9e. Coronary Heart Disease Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	10	5,503	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	7	218	-	-	37	274
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	11	326	9	194
Scott	-	-	-	-	-	-
Sedgwick	102	163	227	252	214	200
Seward	18	434	5	210	-	-
Shawnee	-	-	31	244	4	36
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	5	3,002
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	4	755
Woodson	-	-	-	-	-	-
Wyandotte	43	483	11	144	27	181

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9f. Coronary Heart Disease Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	1,820	367	1,990	275	2,363	263
Frontier	40	268	135	629	117	374
Rural	135	503	110	270	101	240
Dense Rural	452	378	416	220	648	304
Semi-Urban	169	332	113	160	242	248
Urban	1,024	366	1,216	297	1,247	241
Allen	13	1,274	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	4	264	2	210	-	-
Barber	-	-	-	-	-	-
Barton	-	-	30	351	48	457
Bourbon	-	-	-	-	-	-
Brown	-	-	4	455	-	-
Butler	-	-	2	58	36	332
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	45	2,337	-	-
Cheyenne	2	1,651	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	34	4,808	14	1,796
Coffey	8	4,803	-	-	8	1,467
Comanche	-	-	-	-	-	-
Cowley	-	-	33	530	70	622
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	2	1,016
Douglas	5	96	31	301	-	-
Edwards	-	-	-	-	-	-
Elk	-	-	2	1,467	-	-
Ellis	-	-	-	-	6	550
Ellsworth	-	-	-	-	-	-
Finney	61	229	129	262	160	297
Ford	104	447	25	118	132	281
Franklin	-	-	17	700	-	-
Geary	60	857	-	-	11	128
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9f. Coronary Heart Disease Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	8	3,774	-	-
Grant	50	594	56	482	27	274
Gray	-	-	-	-	-	-
Greeley	-	-	12	4,188	-	-
Greenwood	5	1,572	-	-	-	-
Hamilton	-	-	-	-	3	367
Harper	-	-	-	-	-	-
Harvey	42	533	19	217	14	155
Haskell	-	-	7	508	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	20	1,312
Jewell	-	-	-	-	-	-
Johnson	59	159	118	113	204	177
Kearny	2	114	11	348	79	1,811
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	1	36	44	1,734	-	-
Lane	-	-	-	-	-	-
Leavenworth	24	452	49	521	48	336
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	54	451	74	200	22	128
McPherson	20	1,646	4	281	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	42	1,095	7	436
Mitchell	-	-	-	-	-	-
Montgomery	31	821	5	153	-	-
Morris	-	-	-	-	-	-
Morton	6	734	38	2,186	12	469
Nemaha	-	-	-	-	-	-
Neosho	12	566	6	243	52	2,495
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander. Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G9f. Coronary Heart Disease Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	7	2,433
Pawnee	47	4,109	-	-	8	485
Phillips	-	-	-	-	-	-
Pottawatomie	40	2,175	-	-	26	833
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	16	150	-	-	21	111
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	20	759	-	-	47	570
Rooks	-	-	7	2,889	-	-
Rush	-	-	-	-	23	7,212
Russell	-	-	-	-	-	-
Saline	-	-	28	316	106	545
Scott	-	-	-	-	-	-
Sedgwick	374	371	516	347	617	315
Seward	143	802	8	68	81	239
Shawnee	194	471	214	415	162	249
Sheridan	-	-	-	-	-	-
Sherman	25	1,478	9	583	-	-
Smith	-	-	30	12,641	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	18	822	-	-
Stevens	30	954	9	404	37	541
Sumner	-	-	16	358	29	618
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	368	493	288	286	216	203

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10a. Cancer Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	188,291	1,420	191,429	1,325	196,523	1,249
Frontier	9,669	1,384	9,542	1,383	8,688	1,271
Rural	17,661	1,336	18,125	1,313	17,331	1,208
Dense Rural	34,666	1,477	34,031	1,367	34,829	1,324
Semi-Urban	29,638	1,446	31,180	1,402	31,284	1,297
Urban	96,657	1,421	98,466	1,288	104,363	1,222
Allen	1,393	1,829	1,261	1,658	1,388	1,820
Anderson	640	1,455	676	1,596	576	1,131
Atchison	1,298	1,528	1,406	1,552	1,107	1,278
Barber	423	1,398	359	1,059	382	1,312
Barton	2,241	1,610	1,841	1,248	2,170	1,384
Bourbon	1,174	1,480	1,204	1,421	1,030	1,131
Brown	716	1,337	983	1,895	834	1,477
Butler	4,423	1,491	4,450	1,319	4,632	1,244
Chase	370	2,216	401	2,640	305	1,667
Chautauqua	431	1,649	264	849	310	1,106
Cherokee	2,049	1,731	2,288	1,825	2,384	1,935
Cheyenne	222	1,477	210	1,479	156	813
Clark	194	1,588	125	1,054	164	1,291
Clay	547	1,188	472	851	611	1,108
Cloud	730	1,374	759	1,394	807	1,453
Coffey	814	1,796	821	1,556	637	1,139
Comanche	185	1,222	352	3,272	193	1,407
Cowley	3,176	1,720	2,936	1,491	3,042	1,480
Crawford	2,909	1,629	3,130	1,623	2,968	1,468
Decatur	248	1,215	144	619	294	1,384
Dickinson	1,650	1,608	1,588	1,382	1,373	1,160
Doniphan	560	1,290	433	912	648	1,371
Douglas	4,518	1,185	4,917	1,126	5,598	1,119
Edwards	188	945	347	2,098	357	2,263
Elk	439	2,255	283	1,520	240	1,351
Ellis	1,290	999	1,425	1,025	1,315	857
Ellsworth	414	1,111	315	797	380	791
Finney	1,771	1,136	2,050	1,243	2,077	1,163
Ford	2,156	1,585	1,569	1,115	1,460	916
Franklin	1,648	1,316	1,942	1,390	2,130	1,486
Geary	2,142	1,898	1,955	1,696	1,840	1,378
Gove	172	929	199	1,240	206	1,291

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10a. Cancer Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	198	951	257	1,527	244	1,792
Grant	420	1,166	337	900	507	1,304
Gray	338	1,250	464	1,553	282	881
Greeley	12	117	71	826	88	998
Greenwood	734	1,738	933	2,123	529	1,126
Hamilton	296	2,296	155	1,100	119	740
Harper	532	1,437	528	1,367	535	1,352
Harvey	2,359	1,428	2,417	1,389	2,537	1,329
Haskell	115	600	195	882	235	1,028
Hodgeman	183	1,625	164	1,603	157	1,136
Jackson	1,169	1,767	924	1,201	1,149	1,367
Jefferson	1,657	1,637	1,508	1,289	1,642	1,377
Jewell	218	857	244	1,251	250	1,076
Johnson	25,228	1,122	28,713	1,083	28,800	941
Kearny	181	909	231	1,120	311	1,560
Kingman	741	1,626	690	1,482	470	887
Kiowa	150	789	301	1,955	171	1,206
Labette	1,793	1,500	1,945	1,579	2,379	1,959
Lane	248	2,227	163	1,277	78	555
Leavenworth	4,962	1,438	4,387	1,112	5,465	1,240
Lincoln	227	1,163	233	992	317	1,654
Linn	845	1,444	825	1,344	1,022	1,700
Logan	212	1,306	289	1,928	118	617
Lyon	2,049	1,248	2,520	1,488	2,271	1,288
McPherson	1,911	1,282	1,994	1,232	2,179	1,301
Marion	832	1,098	872	1,125	781	1,002
Marshall	692	1,125	675	1,201	680	1,103
Meade	319	1,487	190	788	220	879
Miami	1,948	1,299	2,485	1,431	2,450	1,288
Mitchell	521	1,338	544	1,361	507	1,392
Montgomery	3,303	1,727	3,543	1,869	3,685	1,784
Morris	438	1,298	512	1,307	581	1,478
Morton	172	952	244	1,423	270	1,538
Nemaha	617	1,225	553	1,030	714	1,243
Neosho	1,333	1,479	1,330	1,466	1,251	1,213
Ness	218	1,007	309	1,572	236	1,020
Norton	419	1,368	569	1,694	179	460
Osage	1,154	1,213	1,520	1,652	1,562	1,617

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10a. Cancer Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	467	1,904	380	1,667	308	1,338
Ottawa	594	1,724	517	1,388	394	1,111
Pawnee	576	1,350	559	1,303	718	1,950
Phillips	478	1,415	396	1,163	394	1,194
Pottawatomie	1,391	1,541	1,056	1,033	1,245	1,036
Pratt	810	1,635	591	1,007	730	1,471
Rawlins	176	894	334	2,424	315	2,164
Reno	4,846	1,409	4,882	1,396	5,043	1,356
Republic	382	1,050	208	553	403	1,136
Rice	637	1,152	1,098	2,084	804	1,418
Riley	2,582	1,264	2,136	956	2,068	805
Rooks	501	1,758	403	1,250	418	1,436
Rush	330	1,472	305	1,442	156	608
Russell	533	1,233	614	1,487	532	1,021
Saline	3,478	1,273	4,240	1,439	3,931	1,240
Scott	370	1,366	324	1,270	239	778
Sedgwick	33,883	1,563	34,439	1,430	36,142	1,349
Seward	1,276	1,394	1,267	1,305	1,131	1,102
Shawnee	13,681	1,524	12,834	1,325	13,614	1,312
Sheridan	179	1,043	146	808	200	1,492
Sherman	503	1,530	363	1,078	378	1,093
Smith	374	1,563	324	1,310	284	1,215
Stafford	268	986	299	1,014	232	905
Stanton	113	944	117	1,080	84	664
Stevens	421	1,660	205	772	185	601
Sumner	2,175	1,637	1,966	1,392	2,026	1,441
Thomas	362	943	543	1,236	427	930
Trego	207	1,117	222	1,199	313	2,048
Wabaunsee	528	1,425	606	1,479	382	779
Wallace	148	1,552	40	406	145	2,648
Washington	411	1,143	506	1,539	363	861
Wichita	163	1,159	141	1,119	140	1,110
Wilson	904	1,604	772	1,249	826	1,418
Woodson	284	1,689	396	1,898	596	3,476
Wyandotte	14,385	1,965	13,176	1,736	14,744	1,829

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10b. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	166,811	1,425	165,998	1,324	166,149	1,239
Frontier	9,388	1,400	9,273	1,429	8,226	1,286
Rural	17,231	1,360	17,386	1,326	16,297	1,202
Dense Rural	31,929	1,517	30,854	1,392	31,042	1,356
Semi-Urban	27,475	1,460	27,781	1,377	27,886	1,285
Urban	80,788	1,408	80,640	1,277	82,698	1,194
Allen	1,341	1,821	1,237	1,696	1,356	1,880
Anderson	640	1,477	645	1,573	515	1,019
Atchison	1,257	1,601	1,312	1,537	1,018	1,288
Barber	419	1,418	359	1,079	382	1,361
Barton	2,133	1,667	1,606	1,157	2,056	1,462
Bourbon	1,108	1,463	1,167	1,443	986	1,136
Brown	595	1,260	868	1,871	698	1,357
Butler	4,242	1,493	4,082	1,275	4,232	1,187
Chase	348	2,144	390	2,731	272	1,598
Chautauqua	416	1,673	244	820	303	1,159
Cherokee	1,928	1,716	2,164	1,825	2,287	2,006
Cheyenne	222	1,543	210	1,544	150	810
Clark	194	1,622	125	1,116	164	1,472
Clay	547	1,204	464	849	601	1,116
Cloud	730	1,392	759	1,421	805	1,493
Coffey	786	1,786	786	1,531	631	1,163
Comanche	171	1,132	344	3,339	193	1,436
Cowley	2,753	1,627	2,679	1,507	2,636	1,435
Crawford	2,715	1,588	3,024	1,669	2,875	1,507
Decatur	248	1,222	144	627	294	1,403
Dickinson	1,621	1,641	1,487	1,350	1,316	1,168
Doniphan	560	1,347	420	917	614	1,386
Douglas	4,055	1,185	4,419	1,131	5,087	1,180
Edwards	179	963	347	2,415	332	2,613
Elk	427	2,254	278	1,547	239	1,399
Ellis	1,281	1,016	1,393	1,031	1,279	870
Ellsworth	414	1,179	307	835	349	772
Finney	1,180	1,151	1,438	1,423	1,247	1,207
Ford	1,836	1,875	1,281	1,371	1,120	1,127
Franklin	1,603	1,330	1,698	1,264	2,068	1,537
Geary	1,607	1,993	1,256	1,580	1,225	1,372
Gove	172	934	199	1,266	206	1,327

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10b. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	173	883	240	1,524	188	1,203
Grant	391	1,384	262	966	339	1,276
Gray	338	1,362	370	1,412	262	921
Greeley	12	119	71	856	88	1,085
Greenwood	734	1,778	917	2,166	513	1,138
Hamilton	233	2,043	151	1,197	119	825
Harper	530	1,457	528	1,404	484	1,207
Harvey	2,258	1,500	2,286	1,465	2,236	1,270
Haskell	115	681	170	1,009	162	790
Hodgeman	178	1,628	164	1,692	157	1,190
Jackson	1,075	1,774	852	1,209	976	1,234
Jefferson	1,657	1,677	1,471	1,289	1,608	1,404
Jewell	218	864	244	1,281	220	913
Johnson	23,846	1,151	26,167	1,097	25,738	955
Kearny	141	832	210	1,223	311	2,097
Kingman	730	1,631	690	1,520	470	909
Kiowa	150	800	301	2,030	171	1,274
Labette	1,625	1,479	1,810	1,628	2,154	1,929
Lane	248	2,256	163	1,310	62	461
Leavenworth	4,338	1,455	4,043	1,175	4,689	1,240
Lincoln	227	1,180	233	1,011	317	1,705
Linn	845	1,466	825	1,380	1,015	1,754
Logan	212	1,326	289	2,002	118	638
Lyon	1,749	1,258	2,100	1,437	1,990	1,365
McPherson	1,884	1,300	1,946	1,243	2,166	1,352
Marion	818	1,101	834	1,105	701	928
Marshall	659	1,070	672	1,222	649	1,019
Meade	309	1,610	175	770	214	934
Miami	1,864	1,279	2,310	1,387	2,323	1,277
Mitchell	521	1,345	544	1,376	467	1,351
Montgomery	3,013	1,766	2,794	1,653	3,309	1,809
Morris	438	1,333	466	1,223	581	1,543
Morton	172	1,122	200	1,381	192	1,195
Nemaha	608	1,229	553	1,052	714	1,271
Neosho	1,266	1,467	1,254	1,420	1,196	1,203
Ness	218	1,011	309	1,608	236	1,039
Norton	419	1,480	569	1,790	179	476
Osage	1,143	1,223	1,465	1,625	1,426	1,542

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10b. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	467	1,915	377	1,682	308	1,365
Ottawa	594	1,744	499	1,379	391	1,129
Pawnee	526	1,335	495	1,270	655	2,116
Phillips	478	1,431	396	1,192	389	1,222
Pottawatomie	1,369	1,567	1,031	1,052	1,128	969
Pratt	750	1,576	517	912	579	1,165
Rawlins	176	904	317	2,422	308	2,220
Reno	4,646	1,452	4,433	1,385	4,383	1,289
Republic	382	1,060	208	559	403	1,153
Rice	610	1,165	1,059	2,198	782	1,537
Riley	2,308	1,268	1,919	965	1,758	776
Rooks	501	1,804	380	1,197	418	1,480
Rush	330	1,491	305	1,477	150	597
Russell	533	1,253	608	1,518	532	1,049
Saline	3,219	1,282	3,979	1,515	3,477	1,241
Scott	338	1,328	310	1,370	230	816
Sedgwick	27,550	1,534	27,102	1,396	27,212	1,288
Seward	1,066	1,906	902	1,756	648	1,116
Shawnee	12,014	1,579	10,807	1,338	11,565	1,337
Sheridan	179	1,052	146	815	183	1,462
Sherman	450	1,465	354	1,189	315	936
Smith	374	1,577	324	1,330	284	1,240
Stafford	268	1,016	299	1,056	209	1,014
Stanton	104	988	114	1,477	60	551
Stevens	419	2,082	174	737	147	609
Sumner	2,097	1,651	1,839	1,365	1,835	1,361
Thomas	362	962	543	1,265	427	967
Trego	207	1,126	222	1,227	313	2,103
Wabaunsee	528	1,458	599	1,500	377	790
Wallace	148	1,617	40	421	112	2,002
Washington	411	1,154	506	1,575	363	873
Wichita	163	1,330	88	706	125	1,320
Wilson	892	1,627	764	1,270	809	1,452
Woodson	284	1,748	396	1,942	591	3,625
Wyandotte	8,985	2,051	8,102	1,907	8,407	2,024

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10c. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	13,482	2,121	12,950	1,789	15,168	1,814
Frontier	61	3,422	37	1,549	76	2,725
Rural	63	1,031	79	1,100	78	719
Dense Rural	655	2,146	513	1,480	639	1,592
Semi-Urban	1,265	1,980	1,770	2,459	1,404	1,697
Urban	11,438	2,152	10,534	1,741	12,943	1,864
Allen	44	4,071	24	2,541	13	1,436
Anderson	-	-	-	-	-	-
Atchison	17	374	54	1,301	58	1,220
Barber	-	-	-	-	-	-
Barton	22	1,345	31	2,471	17	1,078
Bourbon	55	2,724	14	606	13	540
Brown	20	2,566	-	-	6	610
Butler	121	3,533	140	4,681	72	1,684
Chase	22	14,832	-	-	-	-
Chautauqua	-	-	-	-	4	2,641
Cherokee	4	419	4	755	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	8	10,566	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	8	6,604	-	-
Cowley	108	2,814	49	1,199	83	2,051
Crawford	-	-	33	1,795	37	1,466
Decatur	-	-	-	-	-	-
Dickinson	-	-	11	2,181	6	683
Doniphan	-	-	-	-	12	1,293
Douglas	352	2,478	166	966	295	1,055
Edwards	9	6,604	-	-	25	37,454
Elk	-	-	-	-	-	-
Ellis	-	-	26	5,783	6	1,981
Ellsworth	-	-	-	-	-	-
Finney	35	3,283	47	2,398	-	-
Ford	25	2,319	38	1,820	-	-
Franklin	45	3,537	40	2,689	-	-
Geary	320	1,892	507	2,262	482	1,911
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10c. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	25	4,480	17	2,032	40	13,551
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	15	756	-	-	35	2,223
Haskell	-	-	-	-	-	-
Hodgeman	5	3,302	-	-	-	-
Jackson	26	10,955	-	-	36	8,317
Jefferson	-	-	-	-	14	3,424
Jewell	-	-	-	-	-	-
Johnson	575	1,368	877	1,180	1,356	1,271
Kearny	-	-	12	8,725	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	160	3,519	89	1,861	126	2,459
Lane	-	-	-	-	-	-
Leavenworth	537	1,879	252	798	524	1,488
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	38	1,668	72	2,574	74	2,108
McPherson	-	-	-	-	11	1,032
Marion	-	-	-	-	-	-
Marshall	-	-	3	1,415	-	-
Meade	-	-	-	-	-	-
Miami	82	3,982	149	6,106	53	1,800
Mitchell	-	-	-	-	-	-
Montgomery	216	2,484	555	6,373	254	2,256
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	14	2,305	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10c. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	43	2,377	37	2,214	24	1,319
Phillips	-	-	-	-	-	-
Pottawatomie	22	3,531	-	-	7	1,778
Pratt	-	-	6	1,165	48	10,231
Rawlins	-	-	-	-	7	23,113
Reno	127	1,936	124	1,629	176	1,979
Republic	-	-	-	-	-	-
Rice	-	-	19	7,535	-	-
Riley	145	1,484	35	501	172	1,135
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	6	2,830	-	-
Saline	194	2,289	187	2,106	123	1,311
Scott	-	-	-	-	-	-
Sedgwick	4,301	2,395	4,011	2,051	5,070	2,203
Seward	85	2,559	54	1,468	129	3,388
Shawnee	1,042	1,656	1,162	1,552	1,320	1,701
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	34	2,898
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	4,631	2,336	4,066	1,947	4,378	2,010

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10d. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	1,594	1,472	1,728	1,439	1,992	1,432
Frontier	42	1,742	75	1,866	22	402
Rural	167	2,403	213	2,346	238	2,296
Dense Rural	541	2,262	369	1,554	580	1,969
Semi-Urban	71	465	292	1,341	288	1,258
Urban	773	1,324	779	1,264	864	1,224
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	15	5,453	34	8,131	31	4,508
Barber	-	-	-	-	-	-
Barton	34	6,646	37	6,738	-	-
Bourbon	-	-	-	-	-	-
Brown	101	3,243	115	3,383	116	3,100
Butler	16	788	18	547	40	1,253
Chase	-	-	11	95,972	-	-
Chautauqua	15	1,657	14	1,342	-	-
Cherokee	101	2,342	55	1,254	39	832
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	10	16,509	16	9,306	-	-
Comanche	14	8,725	-	-	-	-
Cowley	200	5,577	46	1,507	151	3,825
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	5	2,201	-	-	-	-
Doniphan	-	-	11	1,772	6	695
Douglas	107	1,112	65	1,167	99	1,106
Edwards	-	-	-	-	-	-
Elk	-	-	5	3,669	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	12	4,188
Finney	-	-	-	-	22	2,104
Ford	-	-	-	-	28	2,586
Franklin	-	-	128	11,706	18	2,302
Geary	-	-	-	-	19	1,176
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10d. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	23	11,076
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	4	8,805	-	-
Harper	-	-	-	-	22	7,606
Harvey	-	-	11	1,333	12	1,843
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	68	1,776	72	1,552	111	2,168
Jefferson	-	-	29	2,748	-	-
Jewell	-	-	-	-	-	-
Johnson	79	984	85	923	41	366
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	19	939	79	3,635
Lane	-	-	-	-	16	9,971
Leavenworth	10	971	4	307	20	500
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	4	1,761
Logan	-	-	-	-	-	-
Lyon	69	8,131	4	1,321	18	1,309
McPherson	-	-	-	-	-	-
Marion	14	7,185	14	7,704	46	7,441
Marshall	-	-	-	-	-	-
Meade	10	16,509	15	5,453	6	2,476
Miami	-	-	14	1,272	-	-
Mitchell	-	-	-	-	12	4,986
Montgomery	24	385	31	528	44	808
Morris	-	-	16	46,532	-	-
Morton	-	-	3	3,302	-	-
Nemaha	-	-	-	-	-	-
Neosho	35	5,655	4	1,016	50	5,537
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	22	4,686	38	3,822

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10d. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	3	1,415
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	1	220	7	1,216
Pratt	30	40,450	48	30,819	-	-
Rawlins	-	-	-	-	-	-
Reno	13	918	64	2,590	119	4,636
Republic	-	-	-	-	-	-
Rice	-	-	4	1,651	-	-
Riley	-	-	26	6,223	-	-
Rooks	-	-	23	25,843	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	18	1,331	-	-	36	2,163
Scott	-	-	-	-	-	-
Sedgwick	306	1,338	457	1,797	448	1,610
Seward	7	3,556	25	4,322	-	-
Shawnee	167	1,736	94	893	175	1,628
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	3	4,953	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	7	872	10	2,752	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	12	2,935	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	104	1,538	74	959	81	1,090

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10e. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	1,881	836	2,283	808	2,488	710
Frontier	-	-	-	-	10	395
Rural	29	913	18	349	15	313
Dense Rural	164	858	151	610	271	991
Semi-Urban	440	1,587	326	992	271	832
Urban	1,248	732	1,788	817	1,921	676
Allen	8	6,604	-	-	-	-
Anderson	-	-	18	11,217	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	2	1,321	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	18	789	20	959
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	10	3,885
Cloud	-	-	-	-	-	-
Coffey	18	8,725	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	50	2,365	3	190	22	617
Crawford	144	6,129	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	2	629	12	3,173	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	75	703	30	296
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	6	1,101	11	835
Ellsworth	-	-	-	-	-	-
Finney	48	899	25	399	85	1,382
Ford	11	376	9	508	37	964
Franklin	-	-	-	-	-	-
Geary	149	2,284	101	1,294	60	720
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10e. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	70	6,792	11	701	42	2,276
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	5	4,127	-	-
Jewell	-	-	-	-	-	-
Johnson	305	470	481	541	558	504
Kearny	-	-	-	-	-	-
Kingman	11	23,993	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	13	1,400
Lane	-	-	-	-	-	-
Leavenworth	48	802	7	151	61	802
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	8	394	18	636	41	1,607
McPherson	27	5,872	24	3,517	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	31	4,132
Mitchell	-	-	-	-	-	-
Montgomery	-	-	32	3,079	1	132
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	49	9,100	5	3,002
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansas. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10e. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	4	290	63	3,991	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	61	1,251	85	886	70	784
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	12	263	16	215	47	639
Scott	-	-	-	-	-	-
Sedgwick	588	833	995	1,118	777	682
Seward	8	513	-	-	43	1,020
Shawnee	84	878	48	378	61	402
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	10	3,302
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	14	2,841
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	5	3,002
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	223	2,002	182	1,588	434	2,093

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10f. Cancer Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	4,493	765	6,367	794	7,914	766
Frontier	178	860	138	671	195	638
Rural	162	563	300	710	499	944
Dense Rural	1,377	900	1,799	838	1,904	747
Semi-Urban	374	629	628	808	963	857
Urban	2,402	749	3,502	779	4,353	750
Allen	-	-	-	-	3	242
Anderson	-	-	-	-	15	3,966
Atchison	9	594	6	629	-	-
Barber	4	2,401	-	-	-	-
Barton	52	458	167	1,562	72	597
Bourbon	9	2,377	23	4,699	19	2,240
Brown	-	-	-	-	4	472
Butler	44	992	64	794	113	1,108
Chase	-	-	-	-	-	-
Chautauqua	-	-	6	3,048	-	-
Cherokee	16	2,366	13	1,194	14	1,001
Cheyenne	-	-	-	-	6	1,467
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	2	388
Coffey	-	-	19	4,876	6	1,101
Comanche	-	-	-	-	-	-
Cowley	65	1,410	127	1,594	83	912
Crawford	50	730	57	444	31	954
Decatur	-	-	-	-	-	-
Dickinson	22	1,253	32	1,349	48	1,675
Doniphan	-	-	-	-	16	3,674
Douglas	4	77	81	607	80	595
Edwards	-	-	-	-	-	-
Elk	12	3,739	-	-	-	-
Ellis	9	1,143	-	-	19	747
Ellsworth	-	-	8	600	19	1,641
Finney	508	1,073	529	1,027	691	1,018
Ford	284	636	170	377	272	428
Franklin	-	-	31	973	18	636
Geary	66	934	50	1,050	-	-
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10f. Cancer Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	29	350	62	654	145	1,099
Gray	-	-	94	2,603	20	755
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	6	1,132
Hamilton	63	4,159	-	-	-	-
Harper	2	550	-	-	29	3,587
Harvey	16	224	93	882	212	1,424
Haskell	-	-	25	1,001	73	1,326
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	8	1,036
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	423	830	699	719	729	508
Kearny	40	1,160	9	431	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	8	292	16	684	7	353
Lane	-	-	-	-	-	-
Leavenworth	29	306	28	262	139	1,033
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	185	971	310	1,144	148	494
McPherson	-	-	-	-	-	-
Marion	-	-	24	3,596	-	-
Marshall	33	11,925	-	-	31	3,848
Meade	-	-	-	-	-	-
Miami	2	254	2	191	43	1,481
Mitchell	-	-	-	-	15	5,235
Montgomery	50	1,308	75	1,879	72	1,166
Morris	-	-	-	-	-	-
Morton	-	-	41	2,113	78	2,874
Nemaha	-	-	-	-	-	-
Neosho	18	691	1	41	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	11	1,315	33	3,032	30	1,644

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. "-" indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansas. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G10f. Cancer Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	7	889	-	-	32	1,846
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	74	2,299
Pratt	30	2,482	-	-	39	2,033
Rawlins	-	-	17	7,806	-	-
Reno	56	483	172	758	257	1,126
Republic	-	-	-	-	-	-
Rice	27	1,434	16	953	22	1,046
Riley	55	1,142	32	949	25	351
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	6	1,201
Russell	-	-	-	-	-	-
Saline	35	300	52	463	192	903
Scott	32	3,618	14	778	9	772
Sedgwick	1,138	872	1,369	861	1,742	825
Seward	110	713	286	643	295	724
Shawnee	374	845	606	1,054	439	668
Sheridan	-	-	-	-	-	-
Sherman	53	3,134	9	583	37	1,582
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	23	1,000
Stanton	6	508	3	206	24	979
Stevens	2	114	31	475	38	666
Sumner	71	1,325	86	1,429	105	2,712
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	7	1,051	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	53	2,802	15	692
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	434	564	719	643	1,224	921

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: *Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.*

Figure G11a. Diabetes Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	21,155	160	21,930	152	22,985	150
Frontier	788	111	1,098	166	1,012	168
Rural	1,801	137	1,959	137	2,417	184
Dense Rural	4,335	186	4,375	178	3,986	154
Semi-Urban	2,793	137	3,045	138	3,450	150
Urban	11,438	168	11,450	149	12,120	144
Allen	73	94	181	307	184	219
Anderson	15	26	63	131	27	46
Atchison	262	340	144	147	229	235
Barber	51	156	100	307	21	52
Barton	192	133	208	144	316	198
Bourbon	105	137	163	202	152	182
Brown	113	193	138	242	259	502
Butler	462	154	445	135	362	107
Chase	19	104	14	64	15	62
Chautauqua	4	12	10	29	24	75
Cherokee	343	305	369	284	369	270
Cheyenne	4	14	15	73	57	564
Clark	3	17	-	-	23	195
Clay	-	-	43	85	69	130
Cloud	59	98	109	205	58	84
Coffey	58	120	48	89	72	182
Comanche	3	19	-	-	38	357
Cowley	608	317	632	339	440	210
Crawford	323	179	346	204	358	199
Decatur	40	248	13	50	11	39
Dickinson	176	174	46	36	51	42
Doniphan	71	168	143	339	68	204
Douglas	582	152	511	118	545	114
Edwards	36	195	-	-	27	150
Elk	5	17	26	101	28	105
Ellis	219	179	180	135	130	91
Ellsworth	23	58	70	162	105	231
Finney	297	183	268	166	188	115
Ford	133	103	291	211	324	219
Franklin	218	180	212	155	291	216
Geary	230	208	219	190	345	271
Gove	10	39	38	425	39	245
Graham	1	4	16	57	16	71

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. "-" indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11a. Diabetes Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Grant	12	37	41	101	8	23
Gray	31	116	27	89	2	6
Greeley	1	10	4	46	7	88
Greenwood	97	205	53	111	26	46
Hamilton	-	-	44	380	41	282
Harper	58	190	110	278	114	381
Harvey	143	85	185	106	174	84
Haskell	81	340	42	188	45	253
Hodgeman	-	-	16	123	-	-
Jackson	79	114	145	219	22	22
Jefferson	235	239	35	30	93	68
Jewell	44	174	47	168	-	-
Johnson	2,082	93	2,137	81	2,392	79
Kearny	46	232	27	139	83	565
Kingman	60	115	69	139	76	207
Kiowa	34	188	-	-	16	78
Labette	174	134	128	94	243	201
Lane	9	58	19	133	-	-
Leavenworth	485	139	720	184	918	211
Lincoln	48	227	7	32	-	-
Linn	61	91	27	35	69	114
Logan	6	27	20	109	24	115
Lyon	432	257	86	57	170	86
McPherson	203	136	168	94	174	95
Marion	26	31	116	172	82	84
Marshall	68	129	19	28	17	21
Meade	53	248	65	263	70	264
Miami	154	101	119	64	141	82
Mitchell	2	4	40	88	90	300
Montgomery	293	157	461	232	425	230
Morris	50	170	30	69	-	-
Morton	8	40	36	183	35	293
Nemaha	68	124	132	258	117	234
Neosho	217	262	193	215	111	162
Ness	-	-	29	143	2	8
Norton	127	429	30	83	36	91
Osage	42	42	162	172	153	170

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11a. Diabetes Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	47	198	61	248	54	212
Ottawa	49	139	17	42	83	204
Pawnee	69	161	65	139	28	48
Phillips	45	132	47	110	68	178
Pottawatomie	54	66	104	96	76	60
Pratt	92	211	72	118	210	456
Rawlins	31	142	10	41	87	799
Reno	488	148	389	102	718	190
Republic	42	104	65	263	18	41
Rice	52	82	92	149	83	134
Riley	247	129	173	81	150	62
Rooks	48	158	38	108	7	19
Rush	-	-	17	68	-	-
Russell	73	140	62	178	99	301
Saline	235	85	496	166	486	156
Scott	-	-	64	215	68	246
Sedgwick	4,298	198	4,487	187	4,945	186
Seward	164	190	362	390	194	185
Shawnee	1,720	194	1,520	165	1,636	170
Sheridan	47	551	6	30	-	-
Sherman	5	11	218	746	96	361
Smith	1	2	-	-	9	28
Stafford	32	132	80	285	110	390
Stanton	-	-	-	-	-	-
Stevens	62	239	4	15	4	15
Sumner	256	195	367	262	299	209
Thomas	43	119	33	76	121	286
Trego	2	7	4	17	30	160
Wabaunsee	86	230	60	129	117	287
Wallace	24	260	41	421	-	-
Washington	51	157	11	24	33	98
Wichita	29	209	24	181	16	109
Wilson	215	387	162	259	194	416
Woodson	10	32	51	190	45	200
Wyandotte	2,271	310	2,075	267	1,684	214

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11b. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	17,032	147	16,565	132	18,118	140
Frontier	744	110	990	158	838	152
Rural	1,729	138	1,788	132	2,084	164
Dense Rural	3,615	174	3,679	169	3,467	154
Semi-Urban	2,381	127	2,522	126	3,006	148
Urban	8,563	149	7,586	120	8,723	129
Allen	45	58	147	254	184	229
Anderson	15	27	63	133	7	11
Atchison	163	216	102	111	205	228
Barber	51	158	100	314	20	50
Barton	138	98	208	163	268	197
Bourbon	105	143	163	212	152	191
Brown	79	144	99	222	83	139
Butler	407	141	407	127	315	102
Chase	19	108	12	55	15	63
Chautauqua	-	-	10	30	1	3
Cherokee	324	308	362	294	352	276
Cheyenne	4	14	15	74	57	609
Clark	3	17	-	-	23	204
Clay	-	-	43	87	69	133
Cloud	59	99	109	208	58	86
Coffey	58	122	48	91	72	190
Comanche	3	19	-	-	8	45
Cowley	466	265	500	303	325	178
Crawford	323	185	346	216	329	202
Decatur	40	254	13	51	11	40
Dickinson	167	173	39	31	23	17
Doniphan	37	90	118	293	68	216
Douglas	478	139	395	103	429	102
Edwards	27	168	-	-	3	15
Elk	5	18	26	103	28	108
Ellis	219	187	180	141	130	94
Ellsworth	23	60	70	170	105	246
Finney	188	182	170	147	117	105
Ford	102	104	187	208	262	346
Franklin	199	173	202	154	269	216
Geary	138	164	145	189	250	297
Gove	10	39	38	439	39	248

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansas. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11b. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	1	4	16	60	16	75
Grant	12	43	-	-	8	29
Gray	31	121	22	73	2	7
Greeley	1	10	4	47	2	26
Greenwood	97	209	53	114	26	47
Hamilton	-	-	44	735	-	-
Harper	58	194	110	285	113	403
Harvey	140	91	177	115	156	85
Haskell	81	492	21	114	11	43
Hodgeman	-	-	16	127	-	-
Jackson	78	120	67	95	6	7
Jefferson	235	245	35	31	93	70
Jewell	44	175	47	170	-	-
Johnson	1,883	92	1,688	72	2,103	79
Kearny	15	84	27	168	49	507
Kingman	60	117	69	142	76	214
Kiowa	34	190	-	-	16	80
Labette	135	112	113	91	236	220
Lane	9	58	19	136	-	-
Leavenworth	340	115	492	145	769	210
Lincoln	48	229	7	33	-	-
Linn	61	93	27	36	69	119
Logan	6	28	-	-	24	118
Lyon	318	235	77	66	164	104
McPherson	203	139	167	95	174	97
Marion	26	31	116	177	82	86
Marshall	68	130	19	28	17	22
Meade	53	257	65	280	70	289
Miami	151	102	117	65	116	75
Mitchell	2	4	40	88	54	111
Montgomery	230	141	375	222	342	207
Morris	50	175	30	71	-	-
Morton	8	42	36	203	35	413
Nemaha	66	123	132	264	117	241
Neosho	193	244	170	198	71	97
Ness	-	-	29	146	2	8
Norton	127	473	30	85	36	93
Osage	42	42	162	176	153	176

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11b. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	47	199	61	251	54	215
Ottawa	49	140	11	28	60	144
Pawnee	51	134	65	152	28	52
Phillips	45	133	27	64	61	164
Pottawatomie	54	69	99	93	76	63
Pratt	92	222	72	122	210	489
Rawlins	31	144	10	42	87	834
Reno	387	127	295	82	651	189
Republic	42	105	65	266	18	42
Rice	52	86	83	145	83	143
Riley	228	132	100	55	136	65
Rooks	48	161	38	110	7	20
Rush	-	-	17	69	-	-
Russell	71	139	60	181	99	316
Saline	178	68	358	130	442	165
Scott	-	-	36	116	68	271
Sedgwick	3,218	180	2,993	155	3,460	168
Seward	150	242	254	464	145	282
Shawnee	1,340	178	1,022	131	1,056	132
Sheridan	47	562	6	30	-	-
Sherman	5	12	156	652	96	415
Smith	1	2	-	-	9	28
Stafford	32	136	80	301	110	427
Stanton	-	-	-	-	-	-
Stevens	53	259	4	17	-	-
Sumner	253	204	359	269	263	195
Thomas	43	121	33	78	96	231
Trego	2	7	4	17	30	163
Wabaunsee	86	235	60	132	117	298
Wallace	24	270	41	440	-	-
Washington	51	159	11	24	33	100
Wichita	29	244	-	-	-	-
Wilson	208	389	162	266	187	429
Woodson	10	33	51	192	45	204
Wyandotte	1,304	302	996	223	906	230

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansas. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11c. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	2,635	421	3,092	414	2,499	312
Frontier	4	144	-	-	-	-
Rural	20	325	34	469	88	868
Dense Rural	334	971	153	450	161	321
Semi-Urban	220	376	287	443	253	350
Urban	2,057	395	2,618	416	1,997	299
Allen	2	116	32	4,373	-	-
Anderson	-	-	-	-	20	5,816
Atchison	99	3,181	42	974	7	180
Barber	-	-	-	-	-	-
Barton	-	-	-	-	19	1,210
Bourbon	-	-	-	-	-	-
Brown	-	-	14	1,437	-	-
Butler	4	287	38	667	-	-
Chase	-	-	-	-	-	-
Chautauqua	4	6,604	-	-	-	-
Cherokee	15	3,849	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	20	638	32	676	102	1,394
Crawford	-	-	-	-	23	1,017
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	34	5,393	-	-	-	-
Douglas	85	627	42	244	35	167
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	5	236
Ford	13	1,334	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	76	468	49	226	77	290
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11c. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	8	391	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	85	148	236	278	86	94
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	39	858	15	301	-	-
Lane	-	-	-	-	-	-
Leavenworth	106	357	160	453	71	204
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	98	2,534	-	-	6	408
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	3	118	2	77	7	284
Mitchell	-	-	-	-	36	24,392
Montgomery	43	522	68	721	83	1,045
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	23	2,794	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11c. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	18	1,723	-	-	-	-
Phillips	-	-	20	34,899	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	43	590	40	639	17	246
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	4	102	58	534	14	128
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	47	582	24	302	32	327
Scott	-	-	-	-	-	-
Sedgwick	712	422	947	461	838	389
Seward	14	404	9	291	-	-
Shawnee	262	403	391	527	392	531
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	22	1,866
Thomas	-	-	-	-	25	9,631
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	2	777	-	-	7	4,623
Woodson	-	-	-	-	-	-
Wyandotte	807	415	842	398	575	286

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11d. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	391	344	411	338	529	383
Frontier	-	-	19	396	54	1,238
Rural	34	448	25	295	176	1,764
Dense Rural	80	323	129	507	88	291
Semi-Urban	74	342	24	134	17	68
Urban	203	362	214	334	194	279
Allen	26	4,438	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	1	440
Barton	-	-	-	-	8	734
Bourbon	-	-	-	-	-	-
Brown	34	1,101	25	805	176	4,541
Butler	38	1,212	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	23	1,747
Cherokee	-	-	-	-	17	271
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	30	67,417
Cowley	53	1,442	87	2,429	13	383
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	27	4,551
Doniphan	-	-	-	-	-	-
Douglas	19	316	16	184	57	639
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	-	-	-	-	-	-
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11d. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	1	26	33	729	16	248
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	48	419	35	312	18	129
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	7	358
Lane	-	-	-	-	-	-
Leavenworth	39	988	8	614	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	19	15,070	-	-
Lyon	-	-	9	2,972	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	18	273	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: *Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.*

Figure G11d. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	36	1,424	-	-	17	558
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	6	1,101	-	-
Scott	-	-	-	-	-	-
Sedgwick	59	250	105	364	37	121
Seward	-	-	-	-	-	-
Shawnee	28	423	29	346	70	649
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	10	265	21	237	12	161

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11e. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	59	40	131	54	166	47
Frontier	-	-	-	-	-	-
Rural	2	88	-	-	7	146
Dense Rural	-	-	15	93	-	-
Semi-Urban	26	136	-	-	30	79
Urban	31	29	116	58	129	45
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	-	-	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	7	3,082	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	5	317	-	-
Crawford	-	-	-	-	-	-
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	-	-	-	-	24	180
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	-	-
Geary	16	234	-	-	18	163
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11e. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	4	13	42	49	17	15
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	2	43	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	2	1,101	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11e. Diabetes Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	7	2,568
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	-	-
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	-	-	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	10	384	-	-	12	141
Scott	-	-	-	-	-	-
Sedgwick	27	58	72	88	86	76
Seward	-	-	3	126	-	-
Shawnee	-	-	-	-	-	-
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	-	-	-	-	2	18

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11f. Diabetes Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	1,038	217	1,366	207	1,235	136
Frontier	40	277	87	267	96	332
Rural	16	83	112	328	61	112
Dense Rural	306	270	303	210	254	117
Semi-Urban	92	230	212	312	118	138
Urban	584	199	649	171	706	137
Allen	-	-	2	155	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	15	1,246
Barber	-	-	-	-	-	-
Barton	54	779	-	-	21	320
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	13	428	-	-	47	534
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	4	377	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	69	1,733	8	202	-	-
Crawford	-	-	-	-	6	320
Decatur	-	-	-	-	-	-
Dickinson	9	499	-	-	1	53
Doniphan	-	-	-	-	-	-
Douglas	-	-	58	477	-	-
Edwards	9	734	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	109	286	80	256	66	133
Ford	18	121	104	310	62	137
Franklin	19	953	10	475	-	-
Geary	-	-	25	323	-	-
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. "-" indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansas. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11f. Diabetes Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	41	402	-	-
Gray	-	-	5	660	-	-
Greeley	-	-	-	-	5	2,064
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	41	2,018
Harper	-	-	-	-	-	-
Harvey	3	42	-	-	18	130
Haskell	-	-	21	950	34	751
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	62	72	29	42	124	100
Kearny	31	1,196	-	-	34	693
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	-	-	58	352	40	493
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	1	287	-	-
Lyon	16	165	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	18	633
Mitchell	-	-	-	-	-	-
Montgomery	20	667	-	-	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	24	1,668	-	-	40	1,590
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G11f. Diabetes Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	6	3,602	23	2,926
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	5	321	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	22	207	54	342	29	154
Republic	-	-	-	-	-	-
Rice	-	-	9	457	-	-
Riley	15	569	15	307	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	2	777	2	2,641	-	-
Saline	-	-	108	953	-	-
Scott	-	-	28	1,781	-	-
Sedgwick	282	246	251	203	323	172
Seward	-	-	96	310	49	71
Shawnee	90	219	63	129	82	135
Sheridan	-	-	-	-	-	-
Sherman	-	-	62	2,044	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	9	512	-	-	4	180
Sumner	3	92	8	316	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	24	1,300	16	749
Wilson	5	943	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	150	256	190	198	137	100

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: *Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.*

Figure G12a. Unintentional Injury Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	147,833	1,087	147,488	1,076	147,241	1,053
Frontier	10,971	2,064	9,904	2,009	9,301	1,969
Rural	17,862	1,588	16,905	1,594	17,472	1,692
Dense Rural	32,686	1,373	30,250	1,316	31,585	1,397
Semi-Urban	24,646	1,119	25,120	1,135	24,078	1,076
Urban	61,605	846	65,069	859	64,774	819
Allen	978	1,470	1,164	1,950	1,124	1,899
Anderson	792	2,254	682	1,912	578	1,641
Atchison	1,149	1,383	548	730	1,246	1,424
Barber	477	2,340	456	2,016	490	2,545
Barton	1,552	1,147	1,338	1,066	1,707	1,388
Bourbon	1,286	1,858	1,631	2,376	926	1,360
Brown	675	1,383	1,038	2,347	705	1,695
Butler	3,641	1,229	4,140	1,339	3,771	1,215
Chase	255	2,082	243	1,998	121	1,103
Chautauqua	297	1,868	370	2,299	210	1,390
Cherokee	1,481	1,400	1,974	2,013	1,643	1,714
Cheyenne	437	3,675	164	1,521	98	497
Clark	111	1,101	342	3,916	28	229
Clay	282	732	778	2,146	521	1,488
Cloud	640	1,337	420	1,054	740	1,708
Coffey	388	934	389	995	1,084	3,085
Comanche	127	1,891	248	4,187	208	2,456
Cowley	2,609	1,518	2,491	1,451	2,672	1,667
Crawford	2,375	1,180	3,017	1,455	2,345	1,257
Decatur	129	847	286	2,674	230	2,534
Dickinson	1,229	1,484	1,191	1,341	1,399	1,638
Doniphan	546	1,317	281	727	402	955
Douglas	3,841	755	3,838	770	3,872	694
Edwards	652	4,687	115	764	401	3,359
Elk	255	1,663	290	2,529	251	2,072
Ellis	1,272	839	880	617	1,654	1,125
Ellsworth	541	1,833	292	968	447	1,620
Finney	3,158	1,465	2,320	1,232	2,445	1,259
Ford	1,626	941	1,783	1,009	2,432	1,330
Franklin	1,706	1,395	1,702	1,404	1,608	1,320
Geary	1,728	1,021	1,996	1,253	2,488	1,307
Gove	234	2,020	215	1,773	332	2,772

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12a. Unintentional Injury Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	408	4,095	382	3,579	122	1,383
Grant	416	1,063	928	2,328	236	644
Gray	355	1,268	841	2,827	243	877
Greeley	198	3,074	-	-	234	4,598
Greenwood	717	2,206	523	1,741	526	1,885
Hamilton	176	1,309	213	1,687	301	2,434
Harper	546	2,023	816	3,159	616	2,407
Harvey	1,745	1,121	1,518	969	1,653	1,018
Haskell	366	1,774	350	1,758	378	1,859
Hodgeman	153	1,550	73	945	58	797
Jackson	651	1,128	1,200	1,998	540	916
Jefferson	1,722	1,949	1,492	1,711	1,353	1,667
Jewell	205	1,531	141	853	93	911
Johnson	12,117	525	15,995	632	14,428	536
Kearny	737	3,293	380	2,034	520	2,694
Kingman	759	1,966	494	1,340	581	1,723
Kiowa	205	1,580	248	1,656	296	2,645
Labette	1,541	1,488	1,643	1,620	1,989	2,059
Lane	166	1,812	173	2,305	112	1,379
Leavenworth	3,267	928	3,440	960	3,301	856
Lincoln	677	4,893	212	1,631	127	845
Linn	1,428	3,474	720	1,766	599	1,512
Logan	315	2,358	252	2,198	186	1,619
Lyon	2,285	1,252	2,011	1,043	1,985	1,136
McPherson	1,334	890	1,261	926	1,282	933
Marion	1,567	2,572	836	1,572	571	931
Marshall	743	1,558	503	1,132	425	1,021
Meade	434	1,905	333	1,676	375	1,960
Miami	2,220	1,614	1,711	1,151	1,703	1,153
Mitchell	140	479	294	1,056	295	1,144
Montgomery	3,732	2,268	3,576	2,202	2,224	1,456
Morris	333	1,209	506	2,137	300	1,318
Morton	113	732	352	2,297	259	1,882
Nemaha	669	1,460	618	1,469	562	1,265
Neosho	1,697	2,110	943	1,239	1,013	1,402
Ness	280	2,217	215	1,917	490	4,344
Norton	184	670	279	1,049	325	1,247
Osage	1,420	1,924	1,373	1,826	1,495	2,088

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12a. Unintentional Injury Age-Adjusted YPLL* Rates for All Kansans, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	252	1,406	196	1,240	210	1,376
Ottawa	410	1,504	183	537	336	1,401
Pawnee	649	2,104	351	1,133	738	2,393
Phillips	376	1,558	268	1,155	440	1,955
Pottawatomie	1,265	1,415	1,083	1,099	878	857
Pratt	466	973	275	633	950	2,140
Rawlins	84	878	316	3,470	76	789
Reno	3,121	1,014	3,474	1,161	3,839	1,288
Republic	320	1,313	419	2,150	330	1,981
Rice	849	1,617	644	1,561	595	1,318
Riley	1,766	580	1,830	508	2,024	617
Rooks	588	2,451	717	2,943	502	2,328
Rush	189	1,307	277	2,229	279	2,025
Russell	386	1,185	403	1,406	529	1,778
Saline	2,612	989	2,156	782	2,423	928
Scott	113	418	621	2,883	725	3,281
Sedgwick	23,937	1,035	25,813	1,095	26,833	1,079
Seward	1,524	1,251	1,848	1,476	1,562	1,328
Shawnee	8,854	1,057	6,970	821	7,983	934
Sheridan	160	1,199	191	1,823	355	3,915
Sherman	441	1,499	477	1,754	441	1,504
Smith	307	1,769	154	1,000	368	2,538
Stafford	296	1,594	489	2,734	432	2,245
Stanton	390	3,266	410	3,730	79	836
Stevens	416	1,586	787	2,905	505	1,902
Sumner	2,361	1,980	1,795	1,634	1,838	1,670
Thomas	593	1,416	528	1,362	767	1,994
Trego	211	1,429	122	1,066	167	1,325
Wabaunsee	441	1,390	377	1,185	380	1,266
Wallace	192	2,376	129	1,443	142	1,893
Washington	590	2,416	302	1,312	449	1,917
Wichita	103	949	200	2,181	182	1,983
Wilson	1,112	2,422	659	1,550	846	2,161
Woodson	317	1,813	304	2,228	676	5,269
Wyandotte	9,589	1,191	9,013	1,155	8,357	1,062

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12b. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	122,392	1,107	118,051	1,087	115,725	1,080
Frontier	9,697	1,991	8,914	2,024	7,663	1,857
Rural	16,958	1,635	14,741	1,536	15,542	1,709
Dense Rural	26,911	1,395	23,974	1,311	24,888	1,435
Semi-Urban	22,048	1,157	21,648	1,145	20,193	1,083
Urban	46,715	826	48,671	855	47,408	826
Allen	935	1,481	1,072	1,927	1,080	1,983
Anderson	792	2,329	682	1,984	578	1,725
Atchison	1,093	1,470	430	617	1,161	1,450
Barber	424	2,206	410	1,789	490	2,703
Barton	1,086	929	1,261	1,214	1,570	1,562
Bourbon	1,232	1,927	1,436	2,253	896	1,430
Brown	616	1,468	736	1,962	496	1,440
Butler	3,504	1,266	3,818	1,335	3,344	1,188
Chase	255	2,171	243	2,165	63	613
Chautauqua	297	2,036	370	2,509	210	1,518
Cherokee	1,458	1,488	1,634	1,810	1,566	1,826
Cheyenne	379	3,252	128	1,188	98	513
Clark	111	1,179	286	3,692	22	200
Clay	223	586	705	2,011	521	1,568
Cloud	640	1,374	420	1,088	692	1,700
Coffey	335	826	365	999	1,047	3,210
Comanche	127	1,958	203	3,692	208	2,592
Cowley	2,232	1,501	2,172	1,519	2,179	1,688
Crawford	2,218	1,193	2,941	1,562	1,741	1,067
Decatur	129	866	249	2,404	230	2,667
Dickinson	1,229	1,570	1,059	1,266	1,137	1,479
Doniphan	546	1,404	281	775	402	1,082
Douglas	3,103	736	2,998	695	3,476	769
Edwards	596	5,225	115	963	230	2,545
Elk	232	1,571	290	2,651	251	2,201
Ellis	1,218	863	880	664	1,529	1,138
Ellsworth	541	2,098	232	875	447	1,968
Finney	1,014	984	941	1,046	1,301	1,586
Ford	1,099	1,326	925	1,318	807	1,254
Franklin	1,655	1,450	1,484	1,304	1,361	1,194
Geary	962	956	1,144	1,186	1,681	1,425
Gove	234	2,055	215	1,819	281	2,323

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12b. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	332	3,391	264	2,482	89	978
Grant	275	1,142	658	3,279	106	598
Gray	321	1,312	552	2,192	243	1,073
Greeley	99	1,368	-	-	182	4,466
Greenwood	717	2,299	523	1,841	526	2,010
Hamilton	151	1,598	70	581	238	3,134
Harper	546	2,111	802	3,376	428	1,899
Harvey	1,554	1,142	1,435	1,057	1,353	1,003
Haskell	322	2,134	347	2,535	305	2,225
Hodgeman	153	1,627	73	1,009	58	942
Jackson	385	734	932	1,785	362	714
Jefferson	1,660	1,960	1,416	1,707	1,298	1,683
Jewell	205	1,560	141	869	93	930
Johnson	10,679	529	12,928	614	11,990	554
Kearny	207	1,451	341	2,786	355	3,102
Kingman	705	1,868	494	1,412	581	1,825
Kiowa	129	1,022	248	1,738	221	2,172
Labette	1,417	1,566	1,362	1,525	1,820	2,209
Lane	166	1,859	173	2,479	112	1,535
Leavenworth	2,799	982	3,053	1,070	3,004	990
Lincoln	677	4,978	212	1,686	127	873
Linn	1,358	3,426	720	1,834	558	1,470
Logan	315	2,413	252	2,315	186	1,747
Lyon	1,866	1,366	1,394	960	1,549	1,266
McPherson	1,296	915	1,227	954	967	757
Marion	1,567	2,698	730	1,498	529	879
Marshall	743	1,591	503	1,172	385	927
Meade	434	2,249	226	1,391	110	773
Miami	2,208	1,689	1,666	1,182	1,652	1,205
Mitchell	140	488	294	1,091	295	1,194
Montgomery	3,485	2,522	2,894	2,201	2,137	1,774
Morris	333	1,261	506	2,273	300	1,426
Morton	113	897	320	2,821	123	1,224
Nemaha	669	1,491	558	1,370	458	1,068
Neosho	1,564	2,074	887	1,246	927	1,396
Ness	280	2,328	215	2,241	415	4,200
Norton	184	778	240	1,068	325	1,417
Osage	1,392	1,953	1,372	1,898	1,443	2,116

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12b. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic Whites in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	252	1,417	196	1,267	210	1,434
Ottawa	410	1,558	183	549	336	1,478
Pawnee	451	1,679	351	1,339	700	2,744
Phillips	376	1,592	268	1,208	440	2,060
Pottawatomie	1,101	1,289	1,083	1,179	730	763
Pratt	457	1,034	257	654	758	1,928
Rawlins	84	895	316	3,632	76	865
Reno	2,931	1,084	2,894	1,102	3,309	1,300
Republic	320	1,341	360	1,918	330	2,065
Rice	679	1,464	563	1,550	477	1,299
Riley	1,252	523	1,676	591	1,606	633
Rooks	536	2,313	717	3,078	461	2,234
Rush	117	780	277	2,330	279	2,130
Russell	386	1,216	403	1,478	493	1,806
Saline	2,279	1,015	1,696	728	2,009	957
Scott	113	457	466	2,757	703	4,453
Sedgwick	18,718	1,079	19,245	1,123	18,768	1,084
Seward	858	1,690	548	1,314	403	1,061
Shawnee	6,756	1,029	5,438	837	5,893	928
Sheridan	160	1,229	191	1,904	207	2,283
Sherman	345	1,366	477	2,053	318	1,237
Smith	307	1,797	154	1,024	368	2,630
Stafford	296	1,793	404	2,657	338	2,003
Stanton	332	4,059	317	5,145	39	921
Stevens	416	2,255	433	2,310	250	1,517
Sumner	2,230	2,032	1,662	1,654	1,761	1,758
Thomas	593	1,480	528	1,443	566	1,585
Trego	211	1,448	122	1,111	167	1,397
Wabaunsee	428	1,421	335	1,052	346	1,181
Wallace	192	2,678	129	1,585	142	2,186
Washington	590	2,459	246	1,112	392	1,737
Wichita	103	1,316	47	617	140	2,183
Wilson	1,112	2,517	559	1,367	796	2,200
Woodson	317	1,871	245	1,876	661	5,553
Wyandotte	4,660	1,198	5,009	1,444	4,277	1,323

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12c. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	10,051	1,106	9,731	1,090	9,980	1,012
Frontier	76	4,533	118	2,652	51	668
Rural	146	1,088	52	509	163	866
Dense Rural	526	1,073	679	1,557	634	1,199
Semi-Urban	1,266	930	1,405	1,256	1,427	1,274
Urban	8,037	1,124	7,477	1,041	7,705	977
Allen	43	4,940	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	56	737	78	1,889	5	128
Barber	-	-	-	-	-	-
Barton	52	1,741	-	-	-	-
Bourbon	54	1,280	92	4,387	-	-
Brown	-	-	-	-	16	1,485
Butler	55	715	37	650	107	1,575
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	162	20,559	16	2,327
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	59	15,731	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	24	10,787	-	-
Comanche	-	-	-	-	-	-
Cowley	68	1,012	-	-	93	1,630
Crawford	125	2,324	-	-	134	2,485
Decatur	-	-	-	-	-	-
Dickinson	-	-	73	2,844	-	-
Doniphan	-	-	-	-	-	-
Douglas	284	886	478	1,769	99	440
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	23	591
Ford	-	-	-	-	-	-
Franklin	-	-	82	6,432	-	-
Geary	477	1,017	642	1,822	493	1,413
Gove	-	-	-	-	51	176,774

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12c. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	76	52,973	118	33,678	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	180	4,833
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	22	9,270	-	-	17	2,472
Jefferson	62	16,116	53	13,410	-	-
Jewell	-	-	-	-	-	-
Johnson	582	728	1,177	936	1,234	760
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	-	-
Labette	61	1,254	89	1,861	101	1,722
Lane	-	-	-	-	-	-
Leavenworth	396	1,075	264	563	136	315
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	70	1,910	44	1,413	-	-
McPherson	38	3,414	34	3,894	105	2,240
Marion	-	-	28	5,808	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	12	420	-	-	22	852
Mitchell	-	-	-	-	-	-
Montgomery	158	1,483	405	3,365	68	644
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	-	-
Neosho	-	-	-	-	-	-
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12c. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic African Americans in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	74	4,402	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	49	391	149	1,288	178	1,645
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	390	1,235	-	-	119	363
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	90	714	126	1,002
Scott	-	-	-	-	-	-
Sedgwick	2,246	928	2,535	1,112	2,824	1,099
Seward	-	-	31	927	274	7,098
Shawnee	1,375	1,576	732	883	1,017	1,171
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	23	2,154	-	-
Thomas	-	-	-	-	75	14,805
Trego	-	-	-	-	-	-
Wabaunsee	13	7,089	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	57	23,948
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	15	32,718
Wyandotte	3,154	1,350	2,291	1,074	2,395	1,181

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12d. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	1,658	1,159	1,810	1,268	1,770	1,167
Frontier	98	3,716	37	1,144	-	-
Rural	223	2,018	214	2,092	381	3,317
Dense Rural	396	1,327	471	1,601	339	1,052
Semi-Urban	200	821	351	1,577	222	854
Urban	741	1,010	737	948	828	1,099
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	-	-	-	-	38	7,445
Bourbon	-	-	-	-	-	-
Brown	59	1,736	100	2,038	193	3,639
Butler	32	1,020	94	2,873	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	23	544	25	537	61	884
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	-	-	68	1,327	49	1,249
Crawford	32	1,662	-	-	96	3,082
Decatur	-	-	37	46,282	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	363	2,379	115	1,178	38	362
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	-	-	-	-	-	-
Ford	-	-	-	-	-	-
Franklin	-	-	-	-	60	4,251
Geary	51	1,959	-	-	43	1,429
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12d. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	47	35,400	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	14	6,107	63	21,928
Harvey	-	-	46	5,836	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	244	5,743	189	3,715	115	2,112
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	96	1,064	138	1,174	58	489
Kearny	-	-	-	-	-	-
Kingman	54	16,276	-	-	-	-
Kiowa	46	623,636	-	-	-	-
Labette	51	1,976	35	1,405	18	597
Lane	-	-	-	-	-	-
Leavenworth	-	-	58	1,715	29	845
Lincoln	-	-	-	-	-	-
Linn	70	19,595	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	-	-
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	36	539	90	1,145	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	-	-	104	28,917
Neosho	52	3,307	56	6,782	-	-
Ness	-	-	-	-	-	-
Norton	-	-	30	19,262	-	-
Osage	-	-	-	-	52	6,058

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12d. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic American Indian/Alaska Natives in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	-	-	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	-	-	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	49	2,063	-	-	23	747
Republic	-	-	-	-	-	-
Rice	40	14,454	-	-	-	-
Riley	-	-	32	2,107	-	-
Rooks	52	65,542	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	-	-	89	6,395	-	-
Scott	-	-	-	-	-	-
Sedgwick	254	953	308	1,059	451	1,641
Seward	-	-	45	7,923	6	1,651
Shawnee	28	284	118	1,077	218	1,949
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	23	9,691	-	-
Sumner	26	1,654	53	2,752	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	21	3,933
Woodson	-	-	-	-	-	-
Wyandotte	-	-	-	-	34	713

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12e. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	1,684	545	804	216	1,362	312
Frontier	30	2,452	-	-	-	-
Rural	53	1,239	60	1,260	-	-
Dense Rural	398	1,467	1	6	231	658
Semi-Urban	241	594	182	513	205	405
Urban	962	410	561	180	926	273
Allen	-	-	-	-	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	-	-	-	-
Barber	-	-	-	-	-	-
Barton	39	12,684	-	-	-	-
Bourbon	-	-	-	-	-	-
Brown	-	-	-	-	-	-
Butler	-	-	39	1,566	-	-
Chase	-	-	-	-	-	-
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	-	-	-	-
Cheyenne	-	-	-	-	-	-
Clark	-	-	-	-	-	-
Clay	-	-	-	-	-	-
Cloud	-	-	-	-	-	-
Coffey	-	-	-	-	-	-
Comanche	-	-	-	-	-	-
Cowley	64	2,333	-	-	5	241
Crawford	-	-	37	1,829	60	1,477
Decatur	-	-	-	-	-	-
Dickinson	-	-	-	-	-	-
Doniphan	-	-	-	-	-	-
Douglas	55	99	-	-	132	319
Edwards	-	-	-	-	-	-
Elk	-	-	-	-	-	-
Ellis	-	-	-	-	-	-
Ellsworth	-	-	-	-	-	-
Finney	150	2,327	-	-	33	621
Ford	-	-	-	-	-	-
Franklin	50	14,122	-	-	-	-
Geary	139	2,213	-	-	73	875
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12e. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	-	-
Grant	-	-	-	-	-	-
Gray	-	-	-	-	-	-
Greeley	-	-	-	-	-	-
Greenwood	-	-	-	-	-	-
Hamilton	-	-	-	-	-	-
Harper	-	-	-	-	-	-
Harvey	-	-	-	-	-	-
Haskell	-	-	-	-	-	-
Hodgeman	-	-	-	-	-	-
Jackson	-	-	-	-	-	-
Jefferson	-	-	-	-	-	-
Jewell	-	-	-	-	-	-
Johnson	283	334	207	153	150	97
Kearny	-	-	-	-	-	-
Kingman	-	-	-	-	-	-
Kiowa	30	101,126	-	-	-	-
Labette	-	-	-	-	-	-
Lane	-	-	-	-	-	-
Leavenworth	19	396	-	-	-	-
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	-	-	-	-	152	1,820
McPherson	-	-	-	-	-	-
Marion	-	-	-	-	-	-
Marshall	-	-	-	-	-	-
Meade	-	-	-	-	-	-
Miami	-	-	-	-	-	-
Mitchell	-	-	-	-	-	-
Montgomery	-	-	25	2,479	-	-
Morris	-	-	-	-	-	-
Morton	-	-	-	-	-	-
Nemaha	-	-	60	277,292	-	-
Neosho	36	15,822	-	-	41	6,316
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	-	-	1	1,321	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansas. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12e. Unintentional Injury Age-Adjusted YPLL* Rates for Non-Hispanic Asian/Pacific Islanders in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	53	48,988	-	-	-	-
Phillips	-	-	-	-	-	-
Pottawatomie	109	34,104	-	-	-	-
Pratt	-	-	-	-	-	-
Rawlins	-	-	-	-	-	-
Reno	-	-	-	-	14	754
Republic	-	-	-	-	-	-
Rice	-	-	-	-	-	-
Riley	-	-	10	261	-	-
Rooks	-	-	-	-	-	-
Rush	-	-	-	-	-	-
Russell	-	-	-	-	-	-
Saline	52	797	71	1,008	58	958
Scott	-	-	-	-	-	-
Sedgwick	418	449	287	241	356	295
Seward	-	-	-	-	-	-
Shawnee	61	509	-	-	65	439
Sheridan	-	-	-	-	-	-
Sherman	-	-	-	-	-	-
Smith	-	-	-	-	-	-
Stafford	-	-	-	-	-	-
Stanton	-	-	-	-	-	-
Stevens	-	-	-	-	-	-
Sumner	-	-	-	-	-	-
Thomas	-	-	-	-	-	-
Trego	-	-	-	-	-	-
Wabaunsee	-	-	-	-	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	-	-	-	-
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	126	657	67	305	223	919

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12f. Unintentional Injury Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Kansas	11,912	1,002	13,383	918	14,634	866
Frontier	1,070	2,541	736	1,851	1,323	2,672
Rural	482	814	1,341	1,797	1,100	1,363
Dense Rural	4,455	1,334	4,371	1,183	4,819	1,182
Semi-Urban	818	669	1,060	659	1,502	906
Urban	5,087	828	5,803	709	5,890	600
Allen	-	-	92	5,170	-	-
Anderson	-	-	-	-	-	-
Atchison	-	-	40	4,117	-	-
Barber	53	7,987	46	7,995	-	-
Barton	375	2,332	22	257	99	695
Bourbon	-	-	45	3,334	30	6,974
Brown	-	-	96	6,836	-	-
Butler	50	635	74	668	70	575
Chase	-	-	-	-	58	7,116
Chautauqua	-	-	-	-	-	-
Cherokee	-	-	106	8,819	-	-
Cheyenne	58	19,613	36	9,757	-	-
Clark	-	-	56	6,114	-	-
Clay	-	-	73	5,769	-	-
Cloud	-	-	-	-	48	3,828
Coffey	53	5,879	-	-	27	3,371
Comanche	-	-	-	-	-	-
Cowley	245	2,734	225	1,540	133	706
Crawford	-	-	-	-	255	1,851
Decatur	-	-	-	-	-	-
Dickinson	-	-	59	2,216	138	2,344
Doniphan	-	-	-	-	-	-
Douglas	36	293	54	101	45	108
Edwards	56	2,297	-	-	171	6,595
Elk	23	10,337	-	-	-	-
Ellis	54	599	-	-	125	2,158
Ellsworth	-	-	-	-	-	-
Finney	1,994	1,983	1,251	1,489	1,000	1,017
Ford	527	766	823	758	1,591	1,483
Franklin	1	46	58	1,535	135	2,769
Geary	26	431	168	870	198	756
Gove	-	-	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansas. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12f. Unintentional Injury Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Graham	-	-	-	-	33	13,088
Grant	141	809	270	1,423	130	767
Gray	34	1,650	242	4,120	-	-
Greeley	99	9,030	-	-	52	5,113
Greenwood	-	-	-	-	-	-
Hamilton	25	1,793	89	2,587	63	1,757
Harper	-	-	-	-	125	5,536
Harvey	191	1,212	37	307	53	245
Haskell	44	799	3	218	73	1,323
Hodgeman	-	-	-	-	-	-
Jackson	-	-	33	2,519	-	-
Jefferson	-	-	-	-	55	3,218
Jewell	-	-	-	-	-	-
Johnson	477	430	1,146	607	721	293
Kearny	530	6,899	39	710	165	2,713
Kingman	-	-	-	-	-	-
Kiowa	-	-	-	-	75	10,364
Labette	12	748	-	-	50	1,213
Lane	-	-	-	-	-	-
Leavenworth	53	313	65	283	132	495
Lincoln	-	-	-	-	-	-
Linn	-	-	-	-	-	-
Logan	-	-	-	-	-	-
Lyon	349	1,061	495	1,176	284	799
McPherson	-	-	-	-	210	3,915
Marion	-	-	60	2,390	-	-
Marshall	-	-	-	-	40	4,965
Meade	-	-	107	3,654	265	5,915
Miami	-	-	31	1,140	13	457
Mitchell	-	-	-	-	-	-
Montgomery	53	685	134	1,517	19	370
Morris	-	-	-	-	-	-
Morton	-	-	32	1,491	79	2,645
Nemaha	-	-	-	-	-	-
Neosho	45	1,552	-	-	45	1,255
Ness	-	-	-	-	-	-
Norton	-	-	-	-	-	-
Osage	28	4,016	-	-	-	-

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.

Figure G12f. Unintentional Injury Age-Adjusted YPLL* Rates for Hispanics in Kansas, 1999-2013

Geographic Area	1999-2003		2004-2008		2009-2013	
	YPLL	Rate	YPLL	Rate	YPLL	Rate
Osborne	-	-	-	-	-	-
Ottawa	-	-	-	-	-	-
Pawnee	71	3,476	-	-	38	2,297
Phillips	-	-	-	-	-	-
Pottawatomie	55	1,661	-	-	148	2,638
Pratt	9	2,584	18	1,181	192	4,091
Rawlins	-	-	-	-	-	-
Reno	92	499	291	960	232	934
Republic	-	-	-	-	-	-
Rice	130	3,385	79	2,248	65	1,829
Riley	124	894	57	116	297	1,170
Rooks	-	-	-	-	41	14,628
Rush	72	398,282	-	-	-	-
Russell	-	-	-	-	29	4,073
Saline	281	857	210	688	230	1,071
Scott	-	-	127	2,728	22	1,012
Sedgwick	2,301	988	2,560	908	3,047	879
Seward	666	1,036	1,180	1,523	879	1,238
Shawnee	634	889	374	466	592	484
Sheridan	-	-	-	-	148	30,450
Sherman	96	2,468	-	-	50	1,248
Smith	-	-	-	-	-	-
Stafford	-	-	85	3,923	41	1,386
Stanton	58	1,514	93	3,121	40	1,506
Stevens	-	-	331	4,364	203	2,125
Sumner	105	1,440	-	-	32	566
Thomas	-	-	-	-	108	3,645
Trego	-	-	-	-	-	-
Wabaunsee	-	-	42	4,448	-	-
Wallace	-	-	-	-	-	-
Washington	-	-	-	-	-	-
Wichita	-	-	153	6,157	42	1,399
Wilson	-	-	-	-	-	-
Woodson	-	-	-	-	-	-
Wyandotte	1,586	1,011	1,604	761	1,353	602

Note: *Years of Potential Life Lost. See *Technical Notes, Appendix H* for description of Years of Potential Life Lost. “-” indicates that data are unavailable or suppressed. Years of potential life lost are for mortality occurring before age 75. Rates are for three five-year periods and age-adjusted per 100,000 using the U.S. 2000 standard population. Years of potential life lost and rates for Other non-Hispanics are not included as a separate category due to the use of bridged-race estimates for denominators. These counts are included in the count for All Kansans. Using bridged-race estimates, persons identifying as Some Other Race and Two or More Races have been reapportioned to one of the following racial categories: White, African American, American Indian/Alaska Native, and Asian/Pacific Islander.

Source: Kansas Department of Health and Environment, data from combined years 1999-2003, 2004-2008 and 2009-2013.